

# Perspectives on improving photosynthesis to increase crop yield

Roberta Croce,<sup>1,\*</sup> Elizabeth Carmo-Silva,<sup>2</sup> Young B. Cho,<sup>3</sup> Maria Ermakova,<sup>4</sup> Jeremy Harbinson,<sup>5</sup> Tracy Lawson,<sup>6</sup> Alistair J. McCormick,<sup>7,8</sup> Krishna K. Niyogi,<sup>9,10,11,12</sup> Donald R. Ort,<sup>3</sup> Dhruv Patel-Tupper,<sup>9,10</sup> Paolo Pesaresi,<sup>13</sup> Christine Raines,<sup>6</sup> Andreas P.M. Weber,<sup>14</sup> and Xin-Guang Zhu<sup>15</sup>

<sup>1</sup>Department of Physics and Astronomy, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam 1081 HV, the Netherlands

<sup>2</sup>Lancaster Environment Centre, Lancaster University, Lancaster LA1 3SX, UK

<sup>3</sup>Carl R. Woese Institute for Genomic Biology, Department of Plant Biology, University of Illinois, Urbana, IL 61801, USA

<sup>4</sup>School of Biological Sciences, Faculty of Science, Monash University, Melbourne, VIC 3800, Australia

<sup>5</sup>Laboratory of Biophysics, Wageningen University, 6708 WE Wageningen, the Netherlands

<sup>6</sup>School of Life Sciences, University of Essex, Colchester, Essex CO4 3SQ, UK

<sup>7</sup>School of Biological Sciences, Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh EH9 3BF, UK

<sup>8</sup>Centre for Engineering Biology, School of Biological Sciences, University of Edinburgh, Edinburgh EH9 3BF, UK

<sup>9</sup>Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA

<sup>10</sup>Howard Hughes Medical Institute, University of California, Berkeley, CA 94720, USA

<sup>11</sup>Innovative Genomics Institute, University of California, Berkeley, CA 94720, USA

<sup>12</sup>Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>13</sup>Department of Biosciences, University of Milan, 20133 Milan, Italy

<sup>14</sup>Institute of Plant Biochemistry, Cluster of Excellence on Plant Science (CEPLAS), Heinrich Heine University, Düsseldorf 40225, Germany

<sup>15</sup>Key Laboratory of Carbon Capture, Center of Excellence for Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China

\*Author for correspondence: [r.croce@vu.nl](mailto:r.croce@vu.nl)

†Authors are listed alphabetically (with the exception of the lead author/coordinating editor).

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plcell>) is: Roberta Croce ([r.croce@vu.nl](mailto:r.croce@vu.nl)).

## Abstract

Improving photosynthesis, the fundamental process by which plants convert light energy into chemical energy, is a key area of research with great potential for enhancing sustainable agricultural productivity and addressing global food security challenges. This perspective delves into the latest advancements and approaches aimed at optimizing photosynthetic efficiency. Our discussion encompasses the entire process, beginning with light harvesting and its regulation and progressing through the bottleneck of electron transfer. We then delve into the carbon reactions of photosynthesis, focusing on strategies targeting the enzymes of the Calvin–Benson–Bassham (CBB) cycle. Additionally, we explore methods to increase carbon dioxide (CO<sub>2</sub>) concentration near the Rubisco, the enzyme responsible for the first step of CBB cycle, drawing inspiration from various photosynthetic organisms, and conclude this section by examining ways to enhance CO<sub>2</sub> delivery into leaves. Moving beyond individual processes, we discuss two approaches to identifying key targets for photosynthesis improvement: systems modeling and the study of natural variation. Finally, we revisit some of the strategies mentioned above to provide a holistic view of the improvements, analyzing their impact on nitrogen use efficiency and on canopy photosynthesis.

## Introduction

### By Roberta Croce

While photosynthesis is fundamental to life on Earth, powering the growth of plants and sustaining food chains, its solar energy conversion efficiency is surprisingly low, typically below 1% in crops. This low efficiency represents a significant opportunity for improvement, which is especially relevant in the context of a growing global population and pressure on available arable land to climate changes, with the consequent increase in agricultural demand. While substantial progress has been made in enhancing other aspects of crop yield, photosynthesis itself remains a largely untapped area for improvement. This is primarily due to the complexity of the photosynthetic process, which involves a multitude of genes and biochemical pathways. Unlike traits such as disease resistance or plant height, which can often be traced to a few key

genetic changes and can be targeted by traditional breeding, the optimization of photosynthesis demands a deep molecular understanding and precise genetic manipulation. Only recently are the tools and knowledge for such interventions becoming available, making the direct improvement in photosynthesis a relatively new frontier in increasing crop productivity.

Can photosynthesis be improved? Recent proof-of-principle experiments have provided compelling evidence that it can. These studies have explored various innovative approaches, from introducing new biochemical pathways into plant systems to genetic modifications aimed at enhancing light-harvesting or carbon fixation efficiency. For instance, experiments have successfully demonstrated the feasibility of engineering plants to utilize light or assimilate carbon more efficiently or to bypass photorespiratory pathways, thereby boosting overall photosynthetic

Received January 16, 2024. Accepted March 22, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of American Society of Plant Biologists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

efficiency (e.g. Simkin et al. 2015; Kromdijk et al. 2016; South et al. 2019). These ground-breaking experiments show the potential to enhance photosynthesis and open new avenues for increasing crop productivity.

Why, after billions of years of evolution, are there still so many opportunities to enhance photosynthesis? Beyond the common understanding that plants evolve mainly for reproduction and survival rather than maximizing photosynthesis and productivity, there are additional factors. Climate changes, particularly changes in atmospheric carbon dioxide (CO<sub>2</sub>) levels, have made the photosynthetic properties of current plants less suited to modern environments. Enhancing ribulose 1,5-bisphosphate (RuBP) regeneration has been shown to increase photosynthesis in C<sub>3</sub> plants (Lefebvre et al. 2005; Rosenthal et al. 2011), suggesting that the process may not have been fully optimized in response to new environmental conditions. Moreover, inefficiencies in some photosynthetic enzymes and structures may be evolutionary legacies. For example, Rubisco evolved when oxygen (O<sub>2</sub>) was scarce, making differentiation between CO<sub>2</sub> and O<sub>2</sub> less critical. As O<sub>2</sub> became prevalent, increasing the CO<sub>2</sub> specificity of Rubisco meant reducing its catalytic rate, potentially trapping it in a state of low specificity and efficiency (Erb and Zarzycki 2018). These factors suggest that contemporary photosynthesis systems, shaped by environmental changes and evolutionary history, hold potential for optimization and efficiency gains.


This perspective explores a range of promising strategies for improving photosynthetic efficiency. The strategies described here are at various stages of development; some have already demonstrated proof of principle, while others are still in the conceptual phase, collectively representing the state of the art in this field.

## Broadening the spectrum of plants to the far-red

By Roberta Croce

Light is the energy source of photosynthesis. However, only visible photons in the 400 to 700 nm range, the so-called range of photosynthetic active radiation, are used to power photosynthesis in most organisms, including plants. This limits the use of solar energy to <50% of what reaches the Earth's surface (Zhu et al. 2010). For a long time, it has been believed that photons above 700 nm would not carry enough energy to support efficient water oxidation. However, the discovery of cyanobacteria containing pigments absorbing in the far-red (FR) spectral region (Miyashita et al. 1996; Chen et al. 2010) has shown the feasibility of this process. It has thus been proposed that expanding the spectrum of plants up to 750 nm would lead to a gain of light absorption of around 20% (Blankenship and Chen 2013), which is a considerable increase in the energy available for growth. Making use of FR photons would be highly relevant for crops, as plants in the field are close together, and the light reaching the lower leaves is almost exclusively FR (Fig. 1) and currently cannot be used for photosynthesis, resulting in a close-to-zero photosynthetic rate at the bottom of a crop canopy (Srinivasan et al. 2017). The idea of broadening the absorption of crops to the FR is thus considered a promising strategy to improve their productivity (Slattery and Ort 2021; Wu et al. 2023).

How can this be achieved? Although this strategy has not yet been implemented in plants, its success in cyanobacteria serves as an excellent starting point for adapting it to crops. Three main factors are crucial for the efficient use of FR light in the first steps of photosynthesis: (i) the organism's capacity to harvest FR light, (ii) efficient transport of this energy to the photochemical



**Figure 1.** Solar spectrum on the top (yellow) and bottom (brown) of a crop canopy (adapted from Mirkovic et al. 2017). The absorption spectra of three cyanobacteria containing Chl *a* (black), Chl *a* and Chl *f* (red), and Chl *d* (blue) are also shown.

reaction center (RC), and (iii) utilization of this energy for charge separation and stabilization of the charge-separated state. See Elias et al. (2024b) for a recent review.

Plants and algae use chlorophyll *a* (Chl *a*) and Chl *b* and carotenoids to harvest light. These pigments strongly absorb blue and red light but do not absorb in the FR. Some cyanobacteria, however, can synthesize two additional types of chlorophyll, called *d* and *f*, which absorb in the FR region of the spectrum (Gan et al. 2014; Ho et al. 2017). Both these chlorophylls differ from Chl *a* due to the presence of a formyl group in one of the substituents of the tetrapyrrole ring. This also means that a single enzyme is sufficient to transform the red-absorbing Chl *a* into the FR absorbing Chl *d* or *f*. While the enzyme for Chl *d* synthesis has not yet been discovered, the one for Chl *f* is known (Ho et al. 2016) and has been successfully heterologously expressed in cyanobacteria strains incapable of FR acclimation, showing that it leads to the production of Chl *f* (Shen et al. 2019b). These pigments then need to be coordinated to the photosynthetic proteins to form functional pigment-protein complexes. In FR light, alongside the Chl *f* synthase, cyanobacteria express paralogs of some/most photosynthetic proteins involved in pigment binding (Gan et al. 2014). These novel proteins have a higher affinity for Chl *f* in specific sites (Gisriel et al. 2021). However, it has already been shown that even the canonical photosynthetic proteins, such as the components of photosystem I (PSI), can bind Chl *f* (Tros et al. 2020) and that the light-harvesting complexes (LHCs) of plants, which are totally different from those of cyanobacteria, can accommodate both Chl *d* (Elias et al. 2021) and Chl *f* (Elias et al. 2024a).

In addition to the use of Chl *d* and *f*, FR light can also be absorbed by red-shifted Chls *a*. These are present in a small amount in PSI, also of plants, where they are called “red-forms” (Croce and van Amerongen 2013). The FR absorption spectrum of these pigments is due to the strong coupling between two or more Chls *a* (see Slama et al. 2023 for details) which is tuned by the proteins that determine the organization of the pigments. Designing FR Chl *a* binding sites can also be a strategy to enhance the FR absorption of plants without the need for additional Chl types.

After light is absorbed, the energy needs to be efficiently delivered to the RC for charge separation (see Croce and van Amerongen 2020

for more details). This process is made possible by the presence of proteins that organize the pigments. The plant photosynthetic units, PSI and photosystem II (PSII), are modular assemblies where a set of pigment–protein complexes act as an antenna responsible for harvesting light and transferring the excitation energy to the RC. The efficiency of this process hinges on the rate of energy transfer, as faster transfer reduces energy losses. The transfer rate depends on the inter-pigment distance, their mutual orientation, the magnitude of their excited-state transition dipole moments, and their excited-state energy landscapes. With Chl *d* and *f* having larger dipole moments than Chl *a*, and with the organization of the pigments within the complexes remaining largely the same in the studied cyanobacteria complexes (Gisriel et al. 2020, 2023; Hamaguchi et al. 2021), the key factor is the excited-state landscape. The substitution of all Chls *a* with FR absorbing Chls, as in the case of the cyanobacterium *Acaryochloris marina*, does not affect the rate of excitation energy transfer. On the contrary, integrating a few FR chlorophylls into complexes mainly containing Chl *a* can create local energy traps, slowing down the excitation energy transfer rate and reducing efficiency, as observed in cyanobacteria (Mascoli et al. 2020). Careful planning of the number and position of low-energy pigments is therefore crucial to ensure efficient delivery of the absorbed energy to the RC (Mascoli et al. 2022). This would also require the design of modified complexes where the affinity of Chl *d* and *f* is enhanced in specific binding sites.

The presence of low-energy chlorophylls in the RC can influence charge separation efficiency, as these Chls need to perform the same function as Chl *a*, but with less excited-state energy (Nürnberg et al. 2018). This can affect the recombination reactions and, in principle, even increase the possibility of photodamage. The regulation of the midpoint potential of the cofactors in the electron transport chain might be an issue (Viola et al. 2022). This aspect needs to be carefully studied in cyanobacteria to understand how they have overcome possible problems. However, while these factors might be critical for PSII, experimental results indicate that Chl *f*-containing PSI still works very efficiently with Chl *a* in the RC (Tros et al. 2021), suggesting that shifting PSI absorption to the FR might be easier than in the case of PSII.

Finally, the idea is to implement FR absorption in crops using the smart canopy concept, e.g. gradually increasing the absorption of FR photons in the lower part of the canopy (Ort et al. 2015). This allows for better use of light, as the upper leaves in a canopy, typically exposed to full sunlight, would have no advantage in absorbing FR light. In fact, having low-energy Chls in upper canopy leaves could decrease the overall efficiency of charge separation. On the contrary, the absence of FR pigments in the upper leaves allows the penetration of FR light to the bottom of the canopy, where leaves are light-limited and where it will be absorbed and used for photosynthesis. To create this absorption gradient across the canopy, we can again take inspiration from cyanobacteria. More and more species are being discovered that can undergo FR acclimation, meaning that these strains have a normal, Chl *a*-based, photosynthesis apparatus when exposed to white light and only express the Chl *f* synthase and paralogues of the photosynthetic proteins with enhanced ability to bind Chl *f* upon exposure to FR light (Gan et al. 2014; Gisriel et al. 2022). This process is controlled by phytochrome, a photoreceptor that senses the red/FR ratio and triggers the expression of FR-related subunits when needed (Ho et al. 2017). The same control should then be implemented in plants.

In summary, while the utilization of FR light in plant photosynthesis remains to be achieved, the expanding understanding of FR photosynthesis in cyanobacteria is paving the way for this application.

## Enhancing the efficacy of photosynthesis under field conditions: The case for antenna size reduction in crop canopies

By Paolo Pesaresi

In  $C_3$  plant canopies, the photosynthetic machinery saturates at ~25% of maximum solar flux, and this is one of the main factors that constrain productivity in these species (Jansson et al. 2010). Plants probably overinvest in light capture and synthesize high levels of thylakoid antenna proteins and associated chlorophylls because this increases their own fitness by depriving neighboring plants of light and nutrients (Freschet et al. 2011). However, since this trait is found in elite crops, it constitutes a major drawback for monocultures, in which all plants are expected to have the same fitness and yield capacity in order to maximize productivity in cultivated fields (Slattery and Ort 2021).

Reduction of chlorophyll content in leaves could offer a means to mitigate competition for light in monocrop stands, i.e. fields where one crop species is grown at a time, since the level and quality of the light that reaches leaves in the lower canopy should be higher and more uniform, thus boosting overall photosynthetic performance and yields (Blankenship and Chen 2013; Cutolo et al. 2023). Indeed, this strategy has been validated in several studies on cyanobacteria and microalgae (Nakajima and Ueda 1997; Beckmann et al. 2009; Perrine et al. 2012), although the same approach in higher plants has produced discordant results. A decrease in antenna size in tobacco (*Nicotiana tabacum*), for instance, led to an increase of about 25% in above-ground biomass accumulation under high-density cultivation conditions (Kirst et al. 2017). Similarly, beneficial effects were observed in a rice genotype with pale green leaves cultivated under high-light conditions (Gu et al. 2017a). Conversely, the few field studies that have used chlorophyll-deficient soybean (*Glycine max*) mutants showed a marked decrease in leaf mass accumulation and grain yield (Slattery et al. 2017; Sakowska et al. 2018; Genesio et al. 2020). These results underline the complexity of this trait, which is influenced by several factors including the gene/pathway involved, the degree of leaf chlorophyll reduction, the architecture of the crop canopy, as well as the environmental and growth conditions (Cutolo et al. 2023).

Recently, we have used the chemically induced mutant *happy under the sun 1* (*hus1*) to assess the impact of leaf chlorophyll reduction on biomass accumulation and grain yield in barley (Rotasperi et al. 2022). The pale green phenotype of *hus1* is due to a 50% reduction in the chlorophyll content of leaves, which enhances the efficiency of the conversion of light energy into chemical energy without increasing its susceptibility to photoinhibition (Fig. 2). The mutation introduces a premature stop codon in *HvcpSRP43*, which encodes the 43-kDa chloroplast signal recognition particle (cpSRP43) responsible for the delivery of antenna proteins to the thylakoid membranes (Klimyuk et al. 1999), and its truncation resulted in a decrease in the size of photosystem antenna complexes. Furthermore, the *HvcpSRP43* protein has been shown to efficiently chaperone and stabilize glutamyl-tRNA reductase, a rate-limiting enzyme in tetrapyrrole biosynthesis, which enables the insertion of LHCs into thylakoids to be coordinated with chlorophyll biosynthesis (Wang et al. 2018). The dual role of *HvcpSRP43* in chloroplasts makes *hus1* a rather specific mutant, since the reduced antenna size is coupled with a marked reduction in the activity of the tetrapyrrole biosynthetic pathway, which in turn suppresses the formation of highly toxic tetrapyrrole intermediates and pleiotropic photo-oxidative damage. Intriguingly, the *hus1* mutant accumulated biomass and grains at levels comparable to those observed for the control cultivar




**Figure 2.** Phenotypes of the *hus1* mutant and control plants grown under field conditions at Azienda Agraria Sperimentale, Stuard (Parma, Italy). **A**) The *hus1* mutant and the control Sebastian variety at the tillering stage. **B**) *hus1* plants at the heading stage. Courtesy of Lorenzo Genesio (National Research Council, Rome, Italy).

Sebastian, when grown under field conditions at standard density. These findings demonstrate that, when the selective pressure imposed by competition for resources in cultivated fields is attenuated, crops can indeed decrease their investment in antenna proteins and chlorophyll biosynthesis significantly, without detrimental effects on productivity.

Although *cpsrp43* knockout mutants showing the characteristic pale green phenotype have also been reported in other species, including rice (Wang et al. 2016), the *hus1* mutant was chosen for our investigations because a large set of functional genomics tools are available in barley (Rotasperti et al. 2020), making it an optimal model crop in which to test the performance of the several strategies reported to improve photosynthesis efficiency under realistic field conditions (Leister 2023; Smith et al. 2023). For instance, the Fast Identification of Nucleotide variants by droplet Digital PCR technology developed by the Carlsberg Research Laboratory (Knudsen et al. 2022) provides a novel strategy for the rapid identification and isolation of targeted genetic variants in the elite spring barley cultivar RGT Planet. Thanks to this method, mutations that confer advantageous phenotypes on old varieties with low commercial value that carry several other single-nucleotide polymorphisms (SNPs) introduced by chemical mutagenesis—as in the case of the *hus1* mutant isolated within the *HorTILLUS* population (Szurman-Zubrzycka et al. 2018)—can

be transferred to commercially competitive varieties without the need for multiple, time-consuming (several years) backcrosses. Such an approach will make it possible to translate the major gains in biomass and grain yield reported in the last two decades as a consequence of the manipulation of photosystem antenna size in model species and crops, with tests mostly conducted in greenhouses or in small-scale field trials, into yield increases on farms. To this end, collaboration with plant breeders, agronomists, and crop physiologists is needed to select the most appropriate yield-testing protocols, including plot designs that avoid edge effects (which can distort yield estimates), definition of growing plant densities, and standard parameters to define yields (Khaiphob-Burch et al. 2023). Such careful field trials will also allow us to test other major advantages thought to be associated with the pale green phenotype. Independent studies have predicted, for instance, that reductions in chlorophyll content should increase the efficiency of nitrogen use (Song et al. 2017; Sakowska et al. 2018). Similarly, simulations in soybean predict savings of up to 9% of leaf nitrogen upon a 50% reduction in leaf chlorophyll content (Walker et al. 2018). Furthermore, the development of pale green crops and the consequent increase in the fraction of reflected light, i.e. increased albedo, have been shown to mitigate the effects of heat waves triggered by global climate change (Genesio et al. 2021) and improve the efficiency of water



**Figure 3.** Engineering NPQ kinetics via VDE, PsbS, and ZEP overexpression. **A)** Schematic describing the relationships between VDE, PsbS, and ZEP in inducing and relaxing qE and qZ via the interconversion of zeaxanthin (Zea) and violaxanthin (Vio). **B)** Table summarizing differences in published VPZ phenotypes in tobacco (Kromdijk et al. 2016), soybean (De Souza et al. 2022), Arabidopsis (Garcia-Molina and Leister 2020), and potato (Lehretz et al. 2022), based on fluctuating light and field/growth chamber measurements. Triangles describe the directionality of the phenotype relative to wild type (WT). Green triangles indicate hypothesized beneficial photosynthetic efficiency phenotypes, maroon triangles indicate potentially deleterious phenotypes, and yellow lines describe neutral phenotypes. One asterisk (\*) indicates NPQ phenotypes assessed at fluctuations of 500/50  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  rather than 2,000/200  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ . Two asterisks (\*\*) indicate phenotypes reported by Küster et al. (2023). Gray box indicates data not collected in its respective study.

use by reducing canopy temperature (Drewry et al. 2014). This latter aspect is supported by the finding that certain Syrian barley landraces and a few accessions of wild barley (*Hordeum vulgare* spp. *spontaneum*) in Israel, which are adapted to stable and very dry environments, are characterized by pale green leaves (Tardy et al. 1998; Watanabe and Nakada 1999; Galkin et al. 2018).

Overall, the time has come to translate photosynthesis research into the field, using barley as a model crop that can also exploit the availability of large collections of natural genetic diversity (Rotasperi et al. 2020). Numerous publicly funded organizations, such as the “Genomes to Fields Initiative” and the Consultative Group on International Agricultural Research (CGIAR), are conducting field trials that could contribute to achieving this goal with the medium-term objective of transferring this knowledge into other cereals, including wheat, given the high degree of conservation of the photosynthetic machinery in higher plants.

## Accelerating nonphotochemical quenching kinetics to improve photosynthetic efficiency

By Dhruv Patel-Tupper and Krishna K. Niyogi

Photosynthesis needs photoprotection. Excess light can increase the lifetime of the singlet excited state of chlorophyll ( $^1\text{Chl}^*$ ), resulting in

higher yields of the longer-lived ( $\sim\text{ms}$ ) triplet excited state of chlorophyll ( $^3\text{Chl}^*$ ) and photoinhibitory singlet oxygen ( $^1\text{O}_2^*$ ) (Niyogi 1999).  $^3\text{Chl}^*$  and  $^1\text{O}_2^*$  can be quenched by carotenoids, but a ubiquitous first line of defense is the photosystem-scale de-excitation of  $^1\text{Chl}^*$  and dissipation of excess absorbed light energy as heat—a suite of mechanisms that are measured (and referred to) as nonphotochemical quenching (NPQ) of Chl fluorescence (Bassi and Dall’Osto 2021).

NPQ comprises several components defined by their relaxation kinetics and molecular players. In plants, the rapidly induced, energy-dependent quenching (qE) requires the PSII subunit S (PsbS), the xanthophyll zeaxanthin (Zea), and a trans-thylakoid pH gradient ( $\Delta\text{pH}$ ), regulating quenching that is induced and relaxed on a seconds-to-minutes timescale (Bassi and Dall’Osto 2021). A slower type of Zea-dependent but  $\Delta\text{pH}$ -independent quenching (qZ) operates on the timescale of several minutes (Nilkens et al. 2010). Zea is produced by violaxanthin de-epoxidase (VDE) and removed by zeaxanthin epoxidase (ZEP) in a xanthophyll cycle (Yamamoto et al. 1962) (Fig. 3A). The slowest components of NPQ include photoprotective qH (Malnoë et al. 2018) and photoinhibitory qI (Long et al. 1994), which relax on a timescale of hours. In concert, these highly conserved NPQ mechanisms provide the flexibility to cope with light fluctuations at varying intensities and timescales to sustain plant fitness.

Although NPQ protects the photosynthetic apparatus from excess light, it can also compete with and limit photosynthetic efficiency, especially in fluctuating light, where modeling has shown that slow relaxation of NPQ can limit CO<sub>2</sub> assimilation by up to 30% (Zhu et al. 2004). Light intensity fluctuates by time of day, by local shading, and, most dramatically, by the movement of sunflecks in dense crop canopies (Slattery et al. 2018). Optimizing the light-dependent reactions of photosynthesis under dynamic, field-relevant light conditions is a rapidly expanding field with significant agronomic potential (Long et al. 2022; Leister 2023).

A transgenic approach to accelerate the relaxation of NPQ by overexpression of VDE, PsbS, and ZEP (hereafter VPZ, Fig. 3A) increased *N. tabacum* biomass by ~15% (Kromdijk et al. 2016) and elite soybean seed yield by ~20% (De Souza et al. 2022) in small-scale field trials. Although large-scale, multi-location field trials are still needed (Khaipho-Burch et al. 2023), these results point to NPQ as a novel potential target for crop improvement. However, the VPZ approach did not increase biomass in *Arabidopsis thaliana* (Garcia-Molina and Leister 2020) or yield in potato (*Solanum tuberosum*; Lehretz et al. 2022) under greenhouse and simulated fluctuating light conditions.

Here, we consider possible explanations for the varying VPZ results in different plants. Modeling suggests that the stoichiometries of the three proteins are critical (Zaks et al. 2012; De Souza et al. 2022), and thus, it may be necessary to optimize VPZ construct expression for each species depending on native qE and qZ capacities. Plant species with slower NPQ relaxation time constants ( $\tau$ ) and cultivars with higher yield potential may benefit the most from VPZ engineering efforts. Beyond species-specific factors, the literature to date suggests several key phenotypes that may be necessary to achieve greater photosynthetic efficiency by optimizing NPQ.

In all published VPZ studies (tobacco, soybean, *Arabidopsis*, and potato), transformants exhibited faster NPQ relaxation than wild type (WT), when normalized to high-light-acclimated NPQ. However, only in tobacco and two soybean lines (YZ-26-1C and ND-18-44) was faster relaxation also associated with the maintenance of WT high-light NPQ capacity. In other words, maximum NPQ in high light was also increased in VPZ soybean, *Arabidopsis*, and potato lines, which, except for soybean, was associated with decreased biomass relative to WT. The actual magnitude of residual NPQ in low light decreased only in tobacco, YZ-26-1C soybean, and *Arabidopsis* lines. Importantly, tobacco, soybean, and *Arabidopsis* lines showed increases in the low-light effective quantum yield of PSII ( $\Phi$ PSII) (Fig. 3B).

While less residual NPQ and higher  $\Phi$ PSII under fluctuating light are likely strong indicators of improved NPQ kinetics, it is apparent that these traits are not sufficient for increased biomass. Rigorous phenotyping that couple gas exchange measurements of CO<sub>2</sub> assimilation with Chl fluorescence under field-relevant conditions is necessary to demonstrate the impact of faster NPQ kinetics, for example, using 2,000/200  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  fluctuating light assays that maximize fold changes in NPQ while replicating light fluctuations experienced in crop canopies (Slattery et al. 2018; Long et al. 2022). Measuring diurnal de-epoxidation states of the xanthophyll cycle (Kromdijk et al. 2016; De Souza et al. 2022) or the rate of re-epoxidation following HL-to-LL transitions (Küster et al. 2023) may be additional useful proxies for mitigation of the more slowly relaxing qZ component of NPQ. A recent analysis of ZEP-overexpressing lines of the stramenopile alga *Nannochloropsis* (Perin et al. 2023) indicates that for some species, there may also be room to reduce overall NPQ capacity while accelerating NPQ relaxation, highlighting the importance of species-specific phenotypic interrogation.

Natural variation in NPQ may provide nontransgenic avenues to breed for faster NPQ kinetics and higher photosynthetic efficiency. Genome-wide association studies for NPQ have previously revealed diversity in NPQ across rice subspecies and *Arabidopsis* ecotypes (Kasajima et al. 2011; Rungrat et al. 2019). Notably, both studies identified their strongest effect quantitative trait locus within the cis-regulatory regions upstream of the PSBS gene, suggesting NPQ as a trait that is subject to selection. Cowling et al. (2022) measured gas exchange and chlorophyll fluorescence phenotypes across 155 African rice accessions and found that shoot biomass was negatively correlated with NPQ induction capacity and positively correlated with NPQ relaxation rate, consistent with transgenic VPZ phenotypes associated with changes in biomass (Fig. 3B).


Several studies in soybean and maize (*Zea mays*) have identified potential loci for breeding for improved photosynthetic efficiency, using 41 nested association mapping soybean population parents (Wang et al. 2020), 751 diverse maize accessions (Sahay et al. 2023), and 320 multi-parent advanced generation inter-cross maize lines (Ferguson et al. 2023) across multiple years. However, in soybean, there appeared to be little quantitative variation in NPQ relaxation, with photosynthetic differences across lines primarily attributed to differences in Rubisco activation (Wang et al. 2020). Similarly, both maize studies did not resolve robust quantitative variation in residual low-light NPQ specifically, although differences across putative quantitative trait loci in NPQ capacity and maintenance of  $\Phi$ PSII suggest that there may be diverse avenues for fine-tuning light-harvesting efficiency (Ferguson et al. 2023; Sahay et al. 2023). However, crop germplasm screens have not yet revealed differences in NPQ relaxation that are comparable to those in transgenic VPZ plants. Such results could suggest that there are tradeoffs associated with faster NPQ, or more simply, that faster NPQ relaxation has not been selected for in these crops.

Further efforts to improve NPQ kinetics should include accelerating recovery from the slowly reversible types of NPQ, qH, and qI. Nuclear expression of psbA, encoding the D1 subunit of PSII, to minimize qI associated with heat stress is one such approach with demonstrated success across several species (Chen et al. 2020). Additionally, we see tremendous potential in the use of gene editing of endogenous VPZ and other NPQ-related genes in crops to achieve desired NPQ phenotypes (Patel-Tupper et al. 2024). Fine-tuning of VPZ expression and testing diverse VPZ orthologues may be other avenues to bring NPQ-based improvements in photosynthesis closer to their theoretical potential.

## Increasing abundance of the cytochrome *b<sub>6</sub>f* complex to accelerate electron transport rate

By Maria Ermakova

Chloroplast electron transport rate is a primary factor limiting photosynthesis. Classical steady-state quantitative models predict the rate of leaf photosynthesis over a range of intercellular CO<sub>2</sub> partial pressures by considering which biochemical reactions are limiting in certain conditions. In the Farquhar–von Caemmerer–Berry model of C<sub>3</sub> photosynthesis, carbon metabolism (CO<sub>2</sub> availability, Rubisco function) limits assimilation at nonsaturating CO<sub>2</sub> and electron transport limits assimilation under saturating light and saturating CO<sub>2</sub> (intercellular CO<sub>2</sub> partial pressure above 500  $\mu\text{bar}$ ; Farquhar et al. 1980). Based on the model, at current atmospheric CO<sub>2</sub> partial pressure (420 ppm, roughly corresponding to 200  $\mu\text{bar}$  intercellular CO<sub>2</sub>) C<sub>3</sub> photosynthesis is primarily limited by the CO<sub>2</sub> supply to Rubisco, but this limitation is anticipated to



**Figure 4.** Increasing abundance of Cyt  $b_6f$  to accelerate electron transport and enhance the rate of  $C_3$  and  $C_4$  photosynthesis: models' predictions and results from plants overexpressing Rieske FeS subunit of Cyt  $b_6f$  (Rieske-OE). Models' predictions schematically depict simulations obtained with the  $C_3$  photosynthesis model of Farquhar et al. (1980) and  $C_4$  photosynthesis model of von Caemmerer and Furbank (1999). Schematic representations of Rieske-OE results are based on studies conducted in model  $C_3$  plant *A. thaliana* (Simkin et al. 2017b), model  $C_4$  plant *S. viridis* (Ermakova et al. 2019), model  $C_3$  crop *N. tabacum* Petit Havana (Heyno et al. 2022), and model  $C_4$  crop *S. bicolor* Tx430 (Ermakova et al. 2023). In model plants, in line with the models' predictions, Rieske-OE stimulates steady-state electron transport, which results in increased  $CO_2$  assimilation rates at high light and nonlimiting  $CO_2$ . In model crops, Rieske-OE provides transient increases of electron transport, which result in enhanced  $CO_2$  assimilation rate only in the  $C_4$  plant.

transition to electron transport when atmospheric  $CO_2$  exceeds 600 ppm. In the biochemical model of  $C_4$  photosynthesis, electron transport, Rubisco function, and regeneration of RuBP and phosphoenolpyruvate potentially co-limit assimilation under saturating light and saturating  $CO_2$  (intercellular  $CO_2$  partial pressure above 150  $\mu$ bar; von Caemmerer and Furbank 1999; von Caemmerer 2000). The vast difference in the saturating levels of  $CO_2$  between the two photosynthetic pathways is due to the metabolic  $C_4$  cycle of  $C_4$  photosynthesis acting as a carbon-concentrating mechanism. The  $C_4$  cycle operates across two cell types, mesophyll and bundle sheath cells (BSCs), and increases  $CO_2$  partial pressure in BSCs, where Rubisco resides, allowing Rubisco to work at maximum carboxylation rate. Crop models based on these metabolic models of photosynthesis predict yield increases for crops engineered with enhanced electron transport capacity up to 5.2% for wheat and 14.3% for maize (Harbinson and Yin 2023; Wu et al. 2023).

Early experiments on plants with genetically reduced electron transport components identified the thylakoid cytochrome (Cyt)  $b_6f$  complex as the key regulator of the electron transport rate (Price et al. 1995, 1998; Yamori et al. 2011). The complex catalyzes plastoquinone oxidation, considered the slowest reaction of electron transport, and combines it with the translocation of protons across the thylakoid membrane, establishing the proton motive force for ATP generation (Tikhonov 2014). A low luminal pH makes proton translocation more difficult and therefore slows down Cyt  $b_6f$  activity—a phenomenon known as photosynthetic control.

Photosynthetic control via Cyt  $b_6f$  plays an important role in photoprotection by limiting electron flow to PSI and thereby matching the output of the light reactions with the rates of ATP or NADPH consumption by carbon metabolism (Malone et al. 2021). Photosynthetic control therefore represents one molecular mechanism coordinating electron transport and carbon metabolism-related limitations of photosynthesis (Johnson and Berry 2021).

Since Cyt  $b_6f$  was identified as a bottleneck of electron transport, efforts have focused on alleviating this limitation. In higher plants, the complex is comprised of seven essential subunits, of which Rieske FeS and PetM are nuclear-encoded (Schöttler et al. 2015). Enhancing the content of the Rieske FeS subunit through overexpression of the *petC* gene has been established as a method for increasing the abundance and activity of Cyt  $b_6f$  in both  $C_3$  and  $C_4$  plants (Simkin et al. 2017b; Ermakova et al. 2019; Heyno et al. 2022). The increased abundance of the complex in transgenic plants has been confirmed through immunodetection of multiple subunits of the complex in leaves and of the whole complex in isolated thylakoids. Elevated Cyt  $f$  activity has also been detected in thylakoid membranes isolated from leaves of plants overexpressing Rieske FeS.

Increasing abundance of Cyt  $b_6f$  in the model  $C_3$  and  $C_4$  plants, *A. thaliana* and *Setaria viridis*, results in faster electron transport rates through PSI and PSII. Furthermore, in line with the models' predictions, a boost of electron transport capacity enables elevated  $CO_2$  assimilation rates in both model species (Fig. 4). These results confirm that Cyt  $b_6f$  is a bottleneck of electron

transport and that electron transport limits the rate of  $C_3$  and  $C_4$  photosynthesis at saturating light and saturating  $CO_2$ . After genetically engineering plants with increased Cyt  $b_6f$  content was identified as a promising route to improve photosynthesis, overexpression of Rieske FeS has been tested in model  $C_3$  and  $C_4$  crops, tobacco (*N. tabacum* cv. Petit Havana) and sorghum (*Sorghum bicolor* Tx430) (Fig. 4). No differences in steady-state  $CO_2$  assimilation rates and only transient increases of electron transport rate (seen as a faster re-oxidation of plastoquinone in the  $Q_A$  binding site of PSII upon increases of irradiance) are detected in tobacco with enhanced Cyt  $b_6f$  abundance (Heyno et al. 2022). While in sorghum, increasing Cyt  $b_6f$  content does not affect the steady-state rates of electron transport and  $CO_2$  assimilation, it does speed up the induction of photosynthesis after a long period of darkness, and stimulates biomass and grain yield in glasshouse conditions (Ermakova et al. 2023). Overall, results of Rieske FeS overexpression in model crops suggest that the steady-state electron transport rate is no longer limited primarily by Cyt  $b_6f$ .

The photosynthesis rate has undergone significant improvement during crop domestication and early breeding (Huang et al. 2022; Theuvsen et al. 2022). Therefore, it is conceivable that artificial selection of the biggest plants with the greatest yield may have unknowingly increased Rieske FeS content and improved the steady-state electron transport capacity in crops. However, overexpression of Rieske FeS in sorghum has still offered higher yield though transient increases of  $CO_2$  assimilation, highlighting the necessity of studying limitations of crop photosynthesis in dynamic light environments (Kaiser et al. 2017; Long et al. 2022). Looking forward, it is also essential to gain a better understanding of electron transport processes, especially in  $C_4$  plants, where electron transport chains of mesophyll and bundle sheath chloroplasts, distinct in their composition and outputs, cooperatively support seamless operation of biochemical pathways across the two cell types (Munekage 2016; Ermakova et al. 2021b). Cyt  $b_6f$  is the first major bottleneck of electron transport to be successfully alleviated using genetic engineering, demonstrating the viability of this approach for crop improvement. Developing models with more detailed and mechanistic description of electron transport will be instrumental for identifying new targets for optimizing electron transport to boost productivity of crops (Johnson et al. 2021; Bellasio and Ermakova 2022).

## Improving the Calvin–Benson–Bassham cycle

By Christine A. Raines

The Calvin–Benson–Bassham (CBB) cycle evolved over 2 billion years ago (Rasmussen et al. 2008) and is arguably the most important pathway on earth, capturing  $CO_2$  from the atmosphere and converting it into organic molecules that are used directly for the synthesis of isoprenoids, sucrose, starch, phenylpropanoids, thiamine, and nucleotides providing the basis for life on our planet. The CBB cycle is the primary biochemical pathway for the fixation of atmospheric  $CO_2$  in over 85% of plants, named  $C_3$  species, as the first stable product of this cycle is a three-carbon compound, glycerate 3-phosphate (Geiger and Servaites 1994; Sharkey 2019). In the 70 years since the CBB cycle was elucidated, it has been shown to be highly conserved across nature from cyanobacteria to the largest of land plants. The CBB cycle involves 11 enzymes and has three stages: carboxylation carried out by Rubisco, reduction, and RuBP regeneration (Fig. 5). Under light-saturating and  $CO_2$ -limiting conditions, Rubisco activity is the


major determinant of the efficiency of carbon fixation via the CBB cycle. However, as atmospheric  $CO_2$  levels rise and the light intensity decreases, this balance shifts, such that both the reductive and regenerative phases of the CBB cycle that catalyze the synthesis of the  $CO_2$  acceptor molecule RuBP become limiting. Improving photosynthesis has been identified as a target to increase crop yield based on theory, modeling, and empirical studies (Zhu et al. 2010; Simkin et al. 2019; Makino 2021; Raines 2022).

The complexity of the genetics and the biochemistry has made the Rubisco enzyme a challenging target for manipulation. Nevertheless, major efforts to improve photosynthesis have focused on improving Rubisco activity through both direct and indirect approaches. Direct approaches include protein engineering, directed evolution, natural variation screening, and manipulation of expression in transgenic plants (Parry et al. 2012; Yoon et al. 2020; Makino 2021; Gionfriddo et al. 2024). The introduction of  $CO_2$ -concentrating mechanisms (CCMs) from algae and cyanobacteria and engineered synthetic pathways to bypass photosynthesis are indirect approaches being taken and are showing some promise (e.g. South et al. 2019; Shen et al. 2019a). This section will focus on reactions in the CBB cycle other than Rubisco; see the section below by Carmo-Silva for a discussion of Rubisco.

In the 1990s, antisense technology demonstrated that Rubisco did not have total control over  $CO_2$  assimilation under all conditions and identified sedoheptulose 1,7-bisphosphatase (SBPase), fructose 1,6-bisphosphate aldolase (FBPA), and transketolase (TK) as promising targets for the improvement in photosynthesis (Stitt and Schulze 1994; Raines 2003). Based on these studies, a transgenic overexpression approach has shown that increasing the levels of SBPase can improve photosynthesis and growth in algae and a number of plant species including: tobacco (in the field and greenhouse), wheat, and Arabidopsis; in contrast, no positive effect was observed in rice (Lefebvre et al. 2005; Simkin et al. 2015; Driever et al. 2017; Suzuki et al. 2019). Furthermore, tomato plants with increased SBPase activity were found to be more chilling tolerant with increased photosynthetic capacity (Ding et al. 2017). Overexpression of FBPA in tobacco also resulted in positive effects on photosynthesis and biomass (Uematsu et al. 2012; Simkin et al. 2015), and in tomato, an increase in seed weight in both optimal and suboptimal temperatures was observed (Cai et al. 2022). Introduction of the bifunctional cyanobacterial CBB cycle enzyme SBPase/FBPase into tobacco plants, lettuce, and soybean (in elevated  $CO_2$ ) has also resulted in improved  $CO_2$  assimilation and growth (Miyagawa et al. 2001; Tamoi et al. 2006; Kohler et al. 2017; Lopez-Calcagno et al. 2020).

Improvements in RuBP regeneration have also been realized through the introduction of additional proteins that function outside of the CBB cycle. Examples of this approach include combining expression of SBPase and FBPA with either ictB (a cyanobacterial membrane protein of unknown function previously shown to improve  $CO_2$  assimilation) or the H subunit of glycine decarboxylase (GDC) system (shown to increase  $CO_2$  fixation possibly through stimulating the photorespiratory cycle and reducing the negative impact of intermediates on the CBB cycle) in tobacco, which resulted in a further improvement in photosynthesis and growth over single-gene manipulations (Simkin et al. 2015, 2017a). Overexpression of either the endogenous SBPase enzyme or the bifunctional cyanobacterial SBPase/FBPase, together with the algal Cyt C6 protein, not only improved photosynthesis and yield but also increased water use efficiency when grown under field conditions (Lopez-Calcagno et al. 2020). A more recent example is the co-overexpression of SBPase with cytosolic FBPA in tobacco plants, resulting in improvements in biomass, plant height, stem diameter, and pod weight





**Figure 5.** The CBB cycle. Energy in the form of ATP and NADPH (dashed lines) needed to drive the CBB cycle is produced in the thylakoid membrane located electron transport chain. The first step in the CBB cycle is carboxylation (green arrow) catalyzed by Rubisco resulting in the formation of 3-PGA. The next two reactions form the reductive phase (purple arrows) and are catalyzed by phosphoglycerate kinase, forming glycerate 1,3-bisphosphate (BPGA) using ATP and glyceraldehyde 3-phosphate dehydrogenase which forms glyceraldehyde-3-phosphate (GAP) consuming NADPH. Triose phosphate isomerase (TPI) catalyzes the production of dihydroxyacetone phosphate and together with GAP enters the regenerative phase of the cycle (black arrows) catalyzed by fructose 1,6-bisphosphate/sedoheptulose 1,7-bisphosphate aldolase (FBPA), forming sedoheptulose 1,7-bisphosphate (S1,7-BP) and fructose 1,6-bisphosphate (F1,6-BP). SBPase and fructose 1,6-bisphosphatase (FBPase) then produce sedoheptulose 7-phosphate (S7-P) and fructose 6-phosphate (F6-P) which are converted to 5C compounds in reactions catalyzed by TK, ribose 5-P isomerase (RPI), and ribulose 5-phosphate epimerase (RPE) resulting in the formation ribulose 5-P (Ru5P). The final step in the cycle is catalyzed by ribulose 5-phosphate kinase producing the CO<sub>2</sub> acceptor molecule RuBP. The products of the CBB cycle are exported to several biosynthetic pathways for the biosynthesis of isoprenoids, starch, sucrose, shikimate, thiamine, and nucleotides. Rubisco has a competing oxygenase reaction, which results in the formation of 2-phosphoglycerate which enters the photorespiratory pathway (red arrows) (adapted from Raines 2003).

(Li et al. 2022), but in contrast co-overexpression of Rubisco with SBPase in rice did not result in an improvement in photosynthesis (Suzuki et al. 2019). Enhanced photosynthetic capacity was observed in plants in which the expression of a group of CBB cycle genes (FBA1, RCA1, FBPA, and PGK1) in response to higher levels of the Brassinazole resistant 1 transcription factor was increased (Yin et al. 2022). This result suggests that simultaneous overexpression of these enzymes may stimulate the CBB cycle.

Importantly as atmospheric CO<sub>2</sub> rises, theoretical models predict that the limitation of carbon assimilation shifts from Rubisco to RuBP regeneration (Long et al. 2004). Therefore, modifications that improve RuBP regeneration are predicted to stimulate photosynthesis and yield under elevated atmospheric CO<sub>2</sub>. This is supported by experimental evidence using plants grown in free-air CO<sub>2</sub> enrichment (FACE) facilities; when grown at 585 ppm CO<sub>2</sub>, transgenic tobacco plants overexpressing SBPase have greater yield increases at elevated CO<sub>2</sub> (Rosenthal et al. 2011). When both CO<sub>2</sub> and temperature are manipulated, transgenic overexpression of cyanobacterial bifunctional FBPA/SBPase in soybean protects against temperature-induced yield loss under elevated CO<sub>2</sub> (Kohler et al. 2017). These results indicate that improving RuBP regeneration is one approach that could be used to mitigate the effects of climate change on yield, and also demonstrate the importance of testing manipulations in food crops under future climate conditions (Raines 2022).

Advances in kinetic flux and multi-scale modeling have provided novel predictions on how to further enhance the CBB cycle, and the application of rapid high-throughput and iterative approaches will be essential to identify the best candidates to achieve improvements to photosynthesis (Benes et al. 2020; Clapero et al. 2024). Synthetic biology may provide a route to build a completely synthetic, more efficient CO<sub>2</sub> fixation pathway to operate in parallel with the endogenous cycle (Erb and Zarzycki 2016; Löwe and Kremling 2021) or to introduce improved enzymes to operate within the existing cycle. At the same time, new approaches enabling the identification of genetic factors and mechanisms involved in regulating the expression of CBB cycle genes will underpin the application of gene editing technologies to modify this pathway.

## Remarkable, integrated, and complex: paths to improving Rubisco in crops

By Elizabete Carmo-Silva

Rubisco is imperfect yet unique and remarkable (Badger and Sharwood 2023). It catalyzes a rather complex set of reactions during the carboxylation of RuBP and can react with O<sub>2</sub> rather than CO<sub>2</sub>, leading to oxygenation of the sugar phosphate substrate. It

is the only carboxylase that uses substrates and products that are part of central plant metabolism and that is compatible with the CBB cycle (reviewed by Prywes et al. 2023b). Consequently, Rubisco is responsible for assimilating carbon from atmospheric CO<sub>2</sub> into sugars, thereby enabling life on Earth. Rubisco's imperfections have been a research subject and a target for improvement since its discovery (Sharkey 2023). Because the enzyme carboxylates RuBP at relatively slow rates, plants invest large amounts of resources (especially nitrogen and carbon skeletons) into making Rubisco in sufficiently high abundance in the leaves of crop plants to support adequate rates of photosynthetic CO<sub>2</sub> assimilation and plant growth (Carmo-Silva et al. 2015). The biogenesis of Rubisco, consisting of synthesis and assembly into the hexadecameric L<sub>8</sub>S<sub>8</sub> form present in plants, is itself dependent on several ancillary proteins (Bracher et al. 2017), adding further demand on agricultural inputs and impacting resource use efficiency. Thus, while adequate abundance of Rubisco is a necessary consideration in strategies aimed at improving photosynthesis, it needs to be balanced with ensuring efficient use of resources to be compatible with sustainable agricultural practices.

Rubisco catalytic diversity investigations led to the suggestion that tradeoffs associated with the complex reaction mechanism limit the scope for Rubisco improvement (Tcherkez et al. 2006; Savir et al. 2010). A recent phylogenetic analysis demonstrated that, despite being among the 1% slowest evolving enzymes, the continual evolution of Rubisco was associated with improved catalytic efficiency and with greater leaf-level CO<sub>2</sub> assimilation and photosynthetic nitrogen use efficiency (NUE) in C<sub>3</sub> plants (Bouvier et al. 2024). Excitingly, a deep mutational scan using a Rubisco-dependent *Escherichia coli* strain revealed that several highly conserved residues of the loop 6, which folds over the catalytic site and is involved in substrate binding and catalysis, can be mutated without impacting catalysis, suggesting it is possible to target combinations of these residues for future Rubisco engineering efforts (Prywes et al. 2023a).

Rubisco catalytic sites are located at the interface between two large subunits (RbcL), and thus, RbcL has been considered the main source of catalytic diversity and the primary target for engineering efforts (Sharwood 2017). However, despite being far from the catalytic site, just a few changes in the Rubisco small subunit (RbcS) sequence can alter Rubisco catalytic properties (Lin et al. 2022). The role of RbcS in determining the abundance of Rubisco in leaves and adjusting catalytic efficiency has been recently reviewed by Mao et al. (2023) who highlighted the rise of RbcS as a target for improvement that is increasingly tractable as engineering approaches become more robust and widely applicable. Replacement of a crop plant Rubisco by a superior version such as that found in rhodophytes (Oh et al. 2023b) is exciting yet challenging given the need to express adequate levels of RbcS, RbcL, and compatible ancillary proteins. A more straightforward solution might be the use of fast-developing gene editing technologies to mutate specific residues required for enhanced catalysis, when we know what these are.

While the abundance and catalytic properties of Rubisco determine the maximum rate of carboxylation for a given leaf, the activity of Rubisco in crops is regulated by interaction with Rubisco activase (Rca), posttranslational modifications, and the chloroplast stroma environment, which can change rapidly in response to dynamic environmental conditions surrounding the leaf (Amaral et al. 2024). Under fluctuating light conditions, for example, photochemical, biochemical (including Rubisco regulation), and/or diffusional limitations can affect the efficiency of photosynthesis depending on the duration, frequency, and intensity of shade and full sun periods (Long et al. 2022).


Rca is a molecular chaperone that couples ATP hydrolysis with conformational remodeling of inhibited Rubisco catalytic sites, releasing sugar phosphate derivatives that occur naturally and bind tightly and unproductively to the enzyme (Bhat et al. 2017b; Mueller-Cajar 2017). Rca is itself regulated by the redox status, ATP, and Mg<sup>2+</sup> availability in the chloroplast stroma, and some Rca isoforms activate Rubisco more efficiently under fluctuating light (Carmo-Silva and Salvucci 2013). The chaperone is thermolabile, and its ability to restore Rubisco activity is impacted at moderately high temperatures. Progress in understanding the mechanism of Rubisco activation by Rca (Hazra et al. 2015; Bhat et al. 2017a; Flecken et al. 2020) and identification of more thermostable and more efficient Rca isoforms suggest that improvement in Rubisco regulation to maximize carboxylation in current and future warmer climates is possible (Qu et al. 2023; Sparrow-Muñoz et al. 2023; Amaral et al. 2024). Significant unknowns exist in our understanding of Rubisco regulation by Rca, PTMs, and the chloroplast environment (Amaral et al. 2024) and the role of sugar phosphate derivatives and their phosphatases (Orr et al. 2023). Addressing these will aid identification of successful strategies for improving Rubisco, for sustainably increasing crop productivity and climate resilience.

Importantly, to maximize impact, any approach to improve photosynthesis should be considered holistically, and stacking of improvements in various reactions and sub-processes of photosynthesis will be required to ensure that other processes do not become limiting. For example, chloroplast electron transport is also sensitive to heat stress and has been shown to co-limit photosynthesis alongside Rubisco activation (Scafaro et al. 2023). Optimizing Rubisco activity in crop plants requires consideration of Rubisco abundance, Rubisco catalytic properties, and regulation of its activity (Fig. 6A), as well as how Rubisco activity interacts and is coordinated with other plant processes (Fig. 6B). The interaction with the regeneration of RuBP in the CBB cycle is most obvious, but abundance and properties of Rubisco should take into consideration integration with central and specialized metabolism, as well as the specific crop canopy architecture, the needs of the leaves at various canopy layers throughout crop development, and the remobilization of N and C skeletons stored in Rubisco into the crop product to be harvested.


## Introducing CCMs into plants

### By Alistair J. McCormick

The growth of many photosynthetic organisms, including most crops, is limited by the slow rate of CO<sub>2</sub> assimilation by Rubisco and competition with O<sub>2</sub> at the active site resulting in energetically wasteful photorespiration (Bauwe 2023). In response to the shortcomings of Rubisco, nearly every photosynthetic clade has evolved CCMs to supply Rubisco with concentrated CO<sub>2</sub> and preferentially drive CO<sub>2</sub> assimilation over photorespiration. CCMs can be broadly divided into two categories based on biophysical or biochemical processes. Biochemical CCMs initially capture CO<sub>2</sub> as an organic metabolite before re-conversion to CO<sub>2</sub> near Rubisco and include eukaryotic species that perform C<sub>4</sub>, C<sub>2</sub>, and crassulacean acid metabolism (CAM) photosynthesis (Lundgren 2020; Schiller and Bräutigam 2021; Furbank et al. 2023). Biophysical CCMs channel or actively pump in inorganic carbon (Ci, i.e. CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>) to increase the intracellular Ci pool and include prokaryotic autotrophs, eukaryotic algae, hornworts, and seagrasses (Capó-Bauçà et al. 2022; He et al. 2023; Lafferty et al. 2023; Nguyen et al. 2023). All cyanobacteria and some chemoautotrophic bacteria sequester Rubisco within proteinaceous shells called carboxysomes, while almost all algae and some hornworts condense Rubisco into a micro-compartment called the pyrenoid. Both carboxysomes and



**Figure 6.** Improving Rubisco activity in crops. **A)** The activity of Rubisco in a crop leaf can be enhanced by increasing the enzyme abundance, improving its catalytic properties, or optimizing its regulation. Increasing the abundance of Rubisco requires more nitrogen and carbon allocation to Rubisco; some versions of Rubisco show faster rates of carboxylation or higher specificity for  $\text{CO}_2$  over  $\text{O}_2$ ; regulation of the enzyme activity can be enhanced by optimizing the interaction with Rca and ensuring the chloroplast stroma environment is favorable for carboxylation. **B)** To achieve Rubisco-driven crop yield improvements requires consideration of the whole plant. The canopy of the crop will determine which strategy is most promising to improve Rubisco and obtain increased yields and climate resilience. Coordination between photosynthetic sub-processes as well as a productive integration with central and specialized metabolism, plant development, and environmental responses is essential to ensure efficient and sustainable agricultural crop production in present and future climates. A, Rca; C, carbon; N, nitrogen; R, Rubisco; Sugar-P, sugar phosphate derivative.



**Figure 7.** Overview of the requirements for introducing biophysical CCMs into plants. **A)** Engineering a functional pyrenoid-based CCM condensate into a MC chloroplast requires condensation of Rubisco with a linker protein, such as EPYC1 in *Chlamydomonas*, into a pyrenoid-like matrix (shown as a green sphere) that is traversed by thylakoid membranes containing a specialized luminal carbonic anhydrase (CAH3, shown in blue) and bestrophin-like bicarbonate ( $\text{HCO}_3^-$ ) channels on the thylakoid membrane (BST1-3, shown in orange). The addition of a diffusion barrier (e.g. a starch sheath shown in yellow) and an algal  $\text{HCO}_3^-$  channel on the chloroplast envelope (LCIA, shown in red) are predicted to increase the efficiency of  $\text{CO}_2$  capture by Rubisco. **B)** A functional carboxysome-based CCM requires the correct assembly of carboxysome shells that encapsulate cyanobacterial Rubisco and a specialized carboxysomal CA (shown as green hexagons), and active bicarbonate ( $\text{HCO}_3^-$ ) transporters (such as SbtA and/or BicA, shown in dark blue) on the chloroplast envelope that elevate stromal  $\text{HCO}_3^-$  concentrations. Chloroplast stromal CA must be removed to prevent the loss of accumulated  $\text{HCO}_3^-$ .

pyrenoids are supplied with  $\text{HCO}_3^-$  that is rapidly dehydrated by localized carbonic anhydrase (CA) activity to facilitate  $\text{CO}_2$  enrichment at the active sites of Rubisco.

The introduction of biochemical or biophysical CCMs into  $\text{C}_3$  crops is considered a high-risk high-gain engineering strategy to enhance crop yields and resilience (Fig. 7). Models have estimated that the theoretical gains in source leaf photosynthetic efficiency range from 30% to 60%, some of the largest improvements predicted for an engineering approach (Long et al. 2019). Although the levels of translation of such improvements into productivity are debated, for example, the importance of also considering sink demand to take full advantage of source enhancements (Paul 2021), a recent conservative model has predicted an 8% increase in wheat yields, specifically for successful introduction of the cyanobacterial CCM (Wu et al. 2023). Nevertheless, re-constituting any functional CCM in a  $\text{C}_3$  plant remains a

complex endeavor, and engineering each CCM type is beset by specific challenges. In general, advances in our mechanistic understanding of different CCMs have proceeded in parallel with ongoing engineering efforts in a design build test learn (DBTL)-like cycle.

Research efforts to introduce biochemical CCMs into  $\text{C}_3$  plants have largely concentrated on engineering the  $\text{C}_4$  pathway. The  $\text{C}_4$  pathway is estimated to increase  $\text{CO}_2$  levels around Rubisco up to 10-fold above ambient (von Caemmerer and Furbank 2016), which provides  $\text{C}_4$  crops (e.g. maize and sugarcane) with increased photosynthetic efficiencies, growth rates, and in some cases higher water use efficiencies compared to  $\text{C}_3$  crops. Much of this work has been driven by the  $\text{C}_4$  rice project (Furbank et al. 2023), which has focused on introducing the specialized anatomical and biochemical traits of the model  $\text{C}_4$  crop maize into rice. Thus,  $\text{C}_4$  engineers have faced two complex challenges: construction of a “Kranz-like”  $\text{C}_4$  leaf anatomy

with reduced vein spacing and increased organization between mesophyll and BSCs, and biochemical re-regulation for appropriate levels of cell-specific protein expression, for example, sequestration of Rubisco in the bundle sheath chloroplasts. Although the factors involved in the development of the Kranz anatomy are still not fully understood, advances in engineering biology have led to exciting progress in reconstituting  $C_4$  biochemistry in rice. This includes the generation and application of leaf cell-specific expression systems, such that several key  $C_4$  proteins can now be co-expressed and appropriately localized in rice from a single construct (Ermakova et al. 2021a). Attention is now also being expanded to the engineering potential of CAM and  $C_2$  pathways (Lundgren 2020; Schiller and Bräutigam 2021). The CAM pathway could facilitate increased drought tolerance, while  $C_2$  engineering offers some of the benefits of the  $C_4$  pathway with fewer anatomical modifications.

The most efficient biophysical CCMs are carboxysome-based CCMs (cCCMs) and pyrenoid-based CCMs (pCCMs), which can enhance  $CO_2$  concentrations around Rubisco by up to 1,000-fold and 40-fold, respectively (Price et al. 2013; Fei et al. 2022). Biophysical CCMs also function within a single cell, so transfer to  $C_3$  leaf mesophyll cells may be potentially simpler compared to introducing the  $C_4$  pathway. cCCMs have been consistently predicted to increase yield gains in  $C_3$  crops (Nguyen et al. 2023), in part because cyanobacteria have the fastest known form I Rubiscos, with carboxylation turnover rates ( $k_{cat}$ ) up to five times higher than  $C_3$  species (Ang et al. 2023). To date, good progress has been made in understanding the components involved in carboxysome assembly and in reconstructing  $\alpha$ - and  $\beta$ -carboxysomes in plants (Borden and Savage 2022; Blikstad et al. 2023). Recently, Chen et al. (2023) reconstructed the  $\alpha$ -carboxysome from the chemoautotroph *Halothiobacillus neapolitanus* in tobacco chloroplasts, which contained active heterologous Rubisco and CA enzymes. However, functional cCCMs in bacteria still require active uptake of  $HCO_3^-$  and CA activity restricted to carboxysome (Price et al. 2013). Thus, testing whether carboxysomes can enhance growth in plants will require the removal of native chloroplastic CA activity and the introduction of functional  $HCO_3^-$  transporters on the chloroplast envelope. Although the latter remains a long-standing challenge, new screening tools to test the functionality of active  $HCO_3^-$  transporters and channels in *planta* could help to make progress in this area (Förster et al. 2023).

pCCMs are exclusively found in eukaryotes and are likely the most globally abundant CCM type. Pyrenoids are much larger than carboxysomes and characterized by highly diverse morphologies (Barrett et al. 2021), suggesting that there are many ways to achieve pyrenoid formation. The pCCM is best understood in the model alga *Chlamydomonas reinhardtii*, whose pyrenoid is characterized by three architectural features: a liquid-like phase-separated matrix of condensed Rubisco, pyrenoid tubules derived from thylakoid membranes that traverse the matrix to supply  $Cl$ , and a sheath of starch around the matrix that acts as a  $CO_2$  diffusion barrier (Ang et al. 2023; He et al. 2023). Modeling predicts that building a functional *Chlamydomonas* pCCM in a  $C_3$  plant chloroplast could increase  $CO_2$  assimilation rates by up to 3-fold (Fei et al. 2022). pCCMs do not offer the high carboxylation efficiencies of cCCMs but might be more readily plant-compatible based on the appropriate localization of pCCM components thus far expressed (Adler et al. 2022). Furthermore, pCCMs do not require the removal of chloroplastic CA activity and do not need active  $HCO_3^-$  transport, at least at ambient  $CO_2$  levels (Fei et al. 2022). To date, Rubisco condensation into a “proto-pyrenoid” matrix has been achieved in *Arabidopsis* (Atkinson et al. 2020), with work ongoing to reconstitute the two remaining architectural features. Excitingly, recent work in diatoms

and hornworts shows that a variety of solutions may exist to achieve a functional pCCM in plants (Lafferty et al. 2023; Nam et al. 2023; Oh et al. 2023a), for example, by employing a pyrenoid protein shell instead of a starch sheath. Future work could involve creative synthetic strategies that draw from aspects of all CCMs, such as prokaryotic  $HCO_3^-$  transporters coupled to pyrenoids, or a biochemical  $C_4$  pathway localized within a single mesophyll cell (MC) where decarboxylation occurs in a modified chloroplastic carboxysome rather than the BS.


## $C_3$ -to- $C_4$ transition and its potential for improving photosynthesis

By Andreas P.M. Weber

In this section, we aim to provide a concise overview of photorespiration as a limitation on photosynthetic efficiency, the evolution of  $C_4$  photosynthesis via  $C_3$ - $C_4$  intermediates as an adaptation to low atmospheric  $CO_2$  conditions, and efforts to introduce  $C_4$  photosynthesis and bypasses of photorespiration into  $C_3$  plants as a means to increase photosynthetic efficiency. The efficiency of light energy conversion to biomass in  $C_3$  plant photosynthesis is limited by the rate of the oxygenation reaction of Rubisco. In the oxygenation reaction, the acceptor molecule of the CBB cycle, RuBP, is oxidized, not carboxylated, and one of the resulting metabolites, 2-phosphoglycolate (2PG), must be rapidly removed and recycled to 3-phosphoglycerate (3PGA) in a complex pathway called photorespiration (Bauwe 2023; Broncano et al. 2023). The recycling of two molecules of 2PG to 3PGA releases one molecule each of ammonium and  $CO_2$ . It also consumes ATP and redox energy (Fig. 8). Walker et al. (2016) estimated that photorespiration reduces the yield of major  $C_3$  crops by 30% or more. Thus, suppression of photorespiration has great potential to increase crop yield (Eisenhut et al. 2019).

Over the past 20 My, more than 65 dicotyledonous and monocotyledonous plant lineages have independently and convergently evolved a more efficient form of photosynthesis called  $C_4$  photosynthesis (Sage et al. 2012). The frequent evolution of  $C_4$  is thought to have been triggered by a decline in atmospheric  $CO_2$  concentrations from >1,000 ppm to about 350 ppm (Sage et al. 2018). The decrease in  $CO_2$  was accompanied by a drying of the atmosphere and consequently increasing aridity in many parts of the world. Warm temperatures, reduced water availability, and lower atmospheric  $CO_2$  are conditions under which the oxygenation reaction of Rubisco is promoted and photosynthetic efficiency decreases.

$C_4$  plants have a biochemical  $CO_2$  concentration mechanism that increases the  $CO_2$  concentration at the Rubisco site, thereby reducing the rate of photorespiration (Leegood 2002). The biochemical  $CO_2$  pump is supported by a characteristic leaf anatomy called the Kranz anatomy (Sedelnikova et al. 2018). Kranz anatomy is characterized by the formation of two concentric layers of photosynthetic cell types around the leaf vasculature. The inner ring, adjacent to the vasculature, consists of BSCs, which have a large cross-sectional area and are densely filled with chloroplasts containing Rubisco and operating the CBB cycle. BSCs usually perform predominantly cyclic photosynthetic electron transport and have low activity of PSII and hence linear electron transport. The outer ring consists of mesophyll cells, which are always associated with BSCs and face the leaf air space. Mesophyll cells typically contain fewer chloroplasts than BSCs, and they contain little or no Rubisco. MC chloroplasts are capable of linear photosynthetic electron transport. The vascular-bundle sheath-mesophyll (V-BSC-M) cell cluster is the basic unit of a  $C_4$  leaf and is repeated, resulting in a repeating V-BSC-M-M-BSC-V pattern throughout the leaf.



**Figure 8.** Schematic representation of C<sub>3</sub>, C<sub>4</sub>, and C<sub>3</sub>-C<sub>4</sub> intermediate photosynthesis. **A**) C<sub>3</sub> photosynthesis and photorespiration. Both mesophyll (M) and BSCs contain a fully functional photorespiratory pathway. **B**) C<sub>4</sub> photosynthesis. The process of photosynthetic carbon assimilation is divided into two cell types, M and BSCs. M cells act as carbon pumps that increase the CO<sub>2</sub> concentration in BSCs. In BSCs, the CBB cycle operates under elevated CO<sub>2</sub> concentration, which reduces the rate of photorespiration [photorespiratory (PR) pathway not shown; the distribution of the PR pathway between cell types in C<sub>4</sub> is likely equivalent to C<sub>3</sub>-C<sub>4</sub> intermediate photosynthesis, as shown in Fig. 1C]. **C**) C<sub>3</sub>-C<sub>4</sub> intermediate photosynthesis. Photorespiration is shared between M and BSCs, with mitochondrial glycine decarboxylation restricted to the BSCs. Glycine decarboxylation in BSCs locally increases the CO<sub>2</sub> concentration and allows for a more efficient carbon assimilation in this cell type. Please note that many details, cofactors, and pathway intermediates are not shown for clarity. Clp, chloroplast; Gly, glycine; Glyc, glycolate; HP, hydroxypyruvate; Mito, mitochondrion Perox, Peroxisome; Pyr, pyruvate; Rubisco, ribulose 1,5-bisphosphate carboxylase/oxygenase; RuBP, ribulose 1,5-bisphosphate; Ser, serine; TP, triose phosphates.

Mesophyll cells serve as the first carbon-fixing cells. CO<sub>2</sub> from the leaf airspace enters the MC and is converted to HCO<sub>3</sub><sup>-</sup> by CA. HCO<sub>3</sub><sup>-</sup> reacts with phosphoenolpyruvate, catalyzed by phosphoenolpyruvate carboxylase (PEPC), to form the C<sub>4</sub> acid oxaloacetate (OAA). OAA is further converted to malate and/or aspartate, which diffuses along their concentration gradient to the BSCs. In the BSCs, malate and/or aspartate is decarboxylated by one of three decarboxylating enzymes (or a combination thereof; Fig. 1B). These are mitochondrial NAD-malic enzyme (NAD-ME), chloroplastic NADP-malic enzyme (NADP-ME), and PEP carboxykinase (PEPCK).

The predominant pathway of decarboxylation defines the C<sub>4</sub> subtype (NAD-ME, NADP-ME, or PEPCK), although it is controversial whether PEPCK is a distinct subtype or a complementary pathway to either NAD-ME or NADP-ME (Wang et al. 2014a). The C<sub>4</sub> carbon pump raises the CO<sub>2</sub> concentration in BSCs to >1,000 ppm, which suppresses the oxygenation reaction of Rubisco to low levels. However, we emphasize that photorespiration is essential in C<sub>4</sub> plants, as indicated by the lethal phenotype of mutants in the C<sub>4</sub> photorespiratory pathway (Zelitch et al. 2009; Levey et al. 2019).

Based on physiological, anatomical, and biochemical data, as well as computational modeling, it has been proposed that  $C_4$  photosynthesis has gradually evolved from the ancestral  $C_3$  state through  $C_3$ – $C_4$  intermediate states (Sage et al. 2018; Schlüter and Weber 2020). The basic concept is that the photorespiratory pathway loses cell autonomy and is instead split between mesophyll and BSCs (Hylton et al. 1988; Sage et al. 2014). The  $CO_2$  (and ammonia) liberation step of photorespiration, catalyzed by GDC in the mitochondria, ceases to function in mesophyll cells and is restricted to BSCs (Hylton et al. 1988; Rawsthorne et al. 1988). Glycine decarboxylation in BSCs locally increases the  $CO_2$  concentration in this cell type, thereby increasing the efficiency of Rubisco in BSC chloroplasts.

It has been estimated that under conditions that promote high rates of photorespiration (high temperature, low leaf-internal  $CO_2$  concentration due to, e.g. low stomatal conductance),  $CO_2$  accumulates in BSCs at two to three times the level in the mesophyll (Bauwe et al. 1987; Keerberg et al. 2014). Glycine decarboxylation by GDC also releases ammonia, which must be refixed to avoid ammonia toxicity and returned to mesophyll cells to maintain nitrogen balance. It has been proposed that carbon skeletons for ammonia shuttling are provided by mesophyll cells in the form of malate, forming a low-level  $C_4$  metabolic cycle (Mallmann et al. 2014). In this scenario, malate is oxidized to OAA by NADP-malate dehydrogenase. OAA is subsequently transaminated to aspartate, and aspartate is then transported back to the mesophyll cells. Additionally, a portion of the malate can undergo decarboxylation, forming pyruvate, which can be transaminated into alanine, providing an alternative nitrogen transport mechanism to the mesophyll cells. Both malate oxidation via NADP-MDH and oxidative decarboxylation through NAD(P)-ME are facilitated by a predominantly oxidized NAD(P) pool (Bräutigam et al. 2018). If PSII activity is lost from BSCs, it would lead to an increase in the oxidized plastid NADP+ pool. This, in turn, would further promote the evolution toward a full  $C_4$  carbon fixation.

Many staple crops, such as rice and wheat, use  $C_3$  photosynthesis. Given the high efficiency of  $C_4$  photosynthesis, it has been proposed that conversion of these crops to  $C_4$  would result in large yield gains (Hibberd et al. 2008). Indeed, the analysis of a rice new plant type (Sheehy et al. 2001) with large panicles and a low number of tillers showed a grain formation efficiency of only 42%, i.e. a substantial number of juvenile spikelets were not converted into mature, filled spikelets. This finding indicated that rice yield is not limited by sink size, but rather by source strength, i.e. the capacity of leaves to produce and export photo-assimilates (Sheehy et al. 2001). Based on these data, it was estimated that rice yield could be doubled if source strength was increased appropriately. To enhance the source strength, the conversion of rice into a  $C_4$  plant was proposed as a strategic approach, with the potential to increase yield by 50%. This proposition has gained support through FACE experiments. When elite rice varieties were cultivated in FACE conditions with 200 ppm  $CO_2$  concentrations higher than the ambient levels, they produced 13% more grain yield (Zhang et al. 2013). This suggests that improving photosynthesis and thus source strength directly contributes to higher yields in rice.

Significant progress has been made in advancing  $C_4$  photosynthesis in rice (Ermakova et al. 2020). Activation of photosynthetic organelles in the rice BS was achieved by constitutive expression of the maize transcription factors GOLDEN2 or GOLDEN2-LIKE (Wang et al. 2017). Furthermore, five minimal  $C_4$  cycle genes (NADP-ME, PEPC, PPK, MDH, and CA) were successfully expressed in rice from a single construct (Ermakova et al. 2021a). The respective gene products were found to be expressed in the correct cell


types and subcellular compartments. [ $^{13}C$ ]- $CO_2$  labeling showed that some of the labeled carbon is directed through PEPC to malate and aspartate. Rapid labeling of citrate indicated movement of the introduced label toward the tricarboxylic acid (TCA) cycle, while no label was detectable in CBB cycle intermediates. The absence of [ $^{13}C$ ] label in the CBB intermediates indicates that either labeled malate does not enter the plastid stroma or that there is little malate decarboxylation via plastidic NADP-ME (Ermakova et al. 2021a). These results suggest that expression of the core  $C_4$  metabolic enzymes alone is unlikely to be sufficient to establish a functional  $C_4$  cycle. Further development of  $C_4$  rice will likely require the co-expression of organellar metabolite transporters, some of which are unknown. Additionally, it is crucial to prevent the diversion of recently fixed carbon into the TCA cycle, and it may be necessary to inhibit photosynthetic linear electron transport in BSCs to achieve the oxidized NADP+ pool required for decarboxylation by NADP-ME (Bräutigam et al. 2018). Recently, a single promoter TALE system has been reported for tissue-specific expression of multiple transgenes in rice (Danila et al. 2022). In this system, a single cell-specific promoter drives the expression of a synthetic designer transcription activator-like effector that can bind to synthetic TALE-activated promoters (Brückner et al. 2015). This means that multiple genes can be expressed from a single cell-specific promoter in the desired cell type. This technological advance overcomes previous limitations associated with the limited choice of cell-specific promoters for transgene expression in rice and will facilitate transgene stacking, which is critical given the substantial number of transgenes required to implement the trait.

Efforts to establish  $C_4$  photosynthesis (or any other  $C_3$  crop) are ongoing and showing significant progress. In the meantime, other approaches to increase photosynthetic carbon gain in rice and other crops have been explored (Smith et al. 2023). The introduction of synthetic bypasses to photorespiration in rice has been associated with yield increases of up to 15%, which is close to what has been observed in FACE experiments (Shen et al. 2019a). Furthermore, it may be sufficient to install a basic  $C_3$ – $C_4$  intermediate photorespiratory carbon pump in rice, as metabolic modeling suggests that  $C_3$ – $C_4$  intermediate photosynthesis increases carbon gain over a wider range of environmental conditions than  $C_4$ , which is most advantageous under high-light and high-temperature conditions (Bellasio and Farquhar 2019).

## Manipulating stomatal features to improve photosynthesis and water use efficiency

By Tracy Lawson

Gaseous exchange between the leaf interior and the external atmosphere is determined by stomatal conductance ( $g_s$ ), and therefore, stomatal regulatory mechanisms play a pivotal role in determining the rates of photosynthetic carbon assimilation ( $A$ ) and the loss of water through transpiration. Both of these physiological processes hold paramount significance for overall plant performance, productivity, and yield. The importance of  $CO_2$  uptake for photosynthesis is self-evident; however, water loss is equally vital for evaporative cooling and the maintenance of optimal leaf temperature to facilitate photosynthesis (Long et al. 2022), as well as transpiration driving uptake of essential nutrients from the soil to the aerial parts of the plant. The regulation of stomatal aperture, which balances  $CO_2$  uptake and water loss, is thus fundamental in determining plant water use efficiency, determined as  $A/E$  or intrinsic water use efficiency ( $W_i$ ) when assessed directly as a function of  $g_s$ , ( $A/g_s$ ) (Lawson et al. 2010). The significant impact stomata have



**Figure 9.** Schematic diagrams. **A)** Illustrating kinetic responses of A and  $g_s$  to a change in light intensity from low (gray shading) to high intensity (white area). Stomatal responses (blue line) are an order of magnitude slower than A (red line). The shaded green area represents lost CO<sub>2</sub> due to diffusional constraints of slow stomatal opening, while the blue shading represents unnecessary water loss as a result of slow stomatal closure. **B)** Illustrating known mechanisms that increase the rapidity of  $g_s$  responses, including smaller stomata (top), dumbbell-shaped GC (middle), and manipulation of ion transport between GC and SC, at both the plasma membrane and the tonoplast (bottom).

on photosynthetic processes has led to increasing recognition of their potential as valuable targets for manipulation, to enhance crop performance, as well as to develop future crops that can tolerate the challenges posed by climate change (Matthew and Lawson 2019).

Stomatal conductance is the product of both anatomical features and biochemical factors, providing several research avenues for exploitation and manipulation for improved performance. Manipulation of stomatal density (SD) through changes in expression of key genes in the stomatal development and/or patterning pathways has clearly illustrated their potential to either increase the rate of carbon assimilation through removing stomatal limitation (Tanaka et al. 2013; Buckley et al. 2020) or enhance the water use efficiency (W<sub>i</sub>; Bertolino et al. 2019). Although there are various approaches to manipulating stomatal numbers (Bertolino et al. 2019), genes within the epidermal patterning factor (EPF) and epidermal patterning factor-like (EPFL) family have been a particular focus (Harrison et al. 2020). Overexpression of EPF1 and 2 has demonstrated that a reduction in SD (and conductance) resulted in improved drought tolerance and increased water use efficiency (Bertolino et al. 2019; Leakey et al. 2019). However, decreased stomatal conductance usually lowers the rate of carbon assimilation; therefore, it is intriguing that reducing SD (by between 46% and 58%) in the key C<sub>3</sub> crops, rice and wheat (Caine et al. 2019; Dunn et al. 2019) did not impose any diffusional constraints on carbon assimilation. Overexpressing EPFL9/STOMAGEN increases SD leading to a greater conductance associated with enhanced A (Tanaka et al. 2013; Sakoda et al. 2020), highlighting the impact of diffusional constraints on carbon assimilation (Lawson et al. 2012). However, it is important to note that while augmenting SD to increase stomatal conductance may appear promising for removing diffusional constraints (Sakoda et al. 2020), increased water loss erodes water use efficiency (Lawson and Blatt 2014). Furthermore, high SD has been linked to stomatal clustering and impaired stomatal kinetics, resulting in reduced stomatal conductance and rates of carbon assimilation (Dow et al. 2014). These effects are attributed to reductions in critical ion channels necessary for guard cell (GC)

osmoregulation and pore opening, such as reduced K<sup>+</sup> channel activity (Papanatsiou et al. 2016). Therefore, while the manipulation of SD presents a promising avenue for modifying stomatal conductance, it is becoming evident that future endeavors should adopt a more holistic approach that considers both anatomical aspects and functional attributes such as stomatal kinetics (Fig. 9).

The close correlation between the rate of carbon assimilation and stomatal conductance (under steady-state conditions) is well established (Wong et al. 1979); however, it is not always constant (Lawson and Morison 2004), and under dynamic conditions, stomatal responses to changing environmental conditions are an order of magnitude slower than the rate of carbon assimilation, leading to a temporal disconnect between these two processes (Lawson and Blatt 2014). Slow stomatal opening has been reported to reduce carbon assimilation by ca. 10% across a range of species, while slow closure leads to unnecessary water loss that can erode water use efficiency by up to 50% (McAusland et al. 2016). Significant species and cultivar differences in the kinetic responses of stomata that depend on anatomy and biochemistry have been demonstrated (McAusland et al. 2016). This research has led to increasing interest in the rapidity of stomatal conductance and GC regulation as novel targets for improving carbon assimilation and water use efficiency (Lawson and Violet-Chabrand 2019). Recently, Papanatsiou et al. (2019) used optogenetics to enhance solute fluxes in GCs, accelerating stomatal opening and closing, and verifying that enhancing stomatal kinetics improved water use efficiency with no carbon penalty. Engineering ion channels involved in K<sup>+</sup> fluxes in the GCs has further demonstrated the potential to increase the kinetic responses of stomatal conductance and improve water use efficiency (Horaruang et al. 2022). GC metabolism also holds the promise of offering numerous innovative targets for manipulating both the rapidity of stomatal responses and the coordination between carbon assimilation and stomatal conductance (see review by Lemonnier and Lawson 2023).

Smaller stomata have been proposed to have faster kinetics (Drake et al. 2013), most likely facilitated by the greater surface area to volume ratio enabling more rapid solute and water

exchange between GCs and the surrounding cells (Hetherington and Woodward 2003). However, to date little is known about the genetic regulation of stomatal size, and no key targets have been identified for manipulation. Stomatal size has been linked to stomata density, with smaller stomata associated with higher density (Drake et al. 2013); therefore, it may not be possible to manipulate size independently of density. The shape of the GCs and cells that surround them (called subsidiary cells) also influence the speeds at which stomata open and close (Franks and Farquhar 2007). Raissig et al. (2017) confirmed the importance of subsidiary cells in *Brachypodium distachyon* mutants lacking these cells, which resulted in slower kinetics and reduced stomatal conductance. These findings have opened up subsidiary cell metabolism and transport as a potential new avenue of research to explore novel targets for increasing stomata kinetics (Gray et al. 2020) (Fig. 9).

While the scope of this perspective may not encompass all recent advancements, it is worth noting that numerous exciting developments in stomatal research could provide new opportunities to improve crop performance in a changing climate. Nonfoliar stomatal behavior is gaining increasing attention due to the significant impact on photosynthesis and crop yield (especially during periods of plant stress). Additionally, current work has demonstrated considerable water loss associated with these nonfoliar tissues, providing possible new targets to improve whole-plant water use (Lawson and Milliken 2022). While most research primarily concentrates on leaf-level stomatal responses, recent studies have unveiled differences in stomatal kinetics between the adaxial and abaxial stomata (Wall et al. 2022). This discovery opens up the potential to independently manipulate these two surfaces, if the genes responsible for the stomatal patterning on each surface can be elucidated. However, if we are to exploit the potential targets highlighted in this perspective we need a greater understanding of GC signal transduction pathways, metabolism, and the mechanisms that coordinate carbon assimilation and stomatal conductance.

## Natural variation in intrinsic yield potential

By Jeremy Harbinson

The model of Z-scheme oxygenic photosynthesis, with the CBB cycle fixing CO<sub>2</sub>, is the dominant photosynthetic process of the biosphere, and as such, it provides most of the energy and biomass that supports life on Earth. The consistent use of a Z-scheme CBB cycle photosynthetic engine means that there is little variation in the basic physiological and biochemical mechanisms of photosynthesis. However, since the colonization of land by the embryophytes in the late Ordovician (Clarke et al. 2011), this basic mechanism has been adapted by evolution to face the challenges of photosynthesizing optimally in the diverse environmental niches found in the terrestrial biome.

Photosynthesis is a complex process that depends on the cooperative activity of many processes so it can be tuned or optimized along many dimensions, for example, resource use efficiencies such as for light, water, and mineral nutrients, responses to temperature, leaf architecture, and longevity, tolerance to abiotic stress, recovery from stress, etc. These axes for adaptation are individually complex; for example, light can vary in myriad ways (e.g. intensity, spectrum, periodicity, fluctuations over differing time scales) during the life of a leaf or plant. This has resulted in the evolution of variation in photosynthetic traits (e.g. Björkman and Holmgren 1963; Flood et al. 2011; Yamori et al. 2014; Faralli and Lawson 2020). Taking the maximum rate of photosynthetic CO<sub>2</sub> fixation per unit leaf area ( $P_{\max}$ , units  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) as an

example, for a plant using the C<sub>3</sub> photosynthesis mechanism it has been estimated that  $P_{\max}$  should be about  $55 \mu\text{mol m}^{-2} \text{s}^{-1}$  at an irradiance of  $2,000 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Nobel 1991). Some C<sub>3</sub> plants (e.g. desert winter annuals) achieve a slightly higher  $P_{\max}$  [60 to  $65 \mu\text{mol m}^{-2} \text{s}^{-1}$  for *Chylismia* (formerly *Camissonia*) *claviformis* (Mooney et al. 1976) and *Palifoxia linearis* (Werk et al. 1983)]. In contrast, a typical C<sub>3</sub> crop plant will have a  $P_{\max}$  of 20 to  $30 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Nobel 1991), deciduous forest tree species a  $P_{\max}$  of about  $10 \mu\text{mol m}^{-2} \text{s}^{-1}$ , while for permanently shaded rainforest understory species,  $P_{\max}$  is 1 to  $5 \mu\text{mol m}^{-2} \text{s}^{-1}$ .

Despite the variation for photosynthesis observed in natural systems, it has been argued that photosynthesis in crop plants cannot be improved because it has already been optimized by natural selection and evolution. This argument is, however, only partly correct because plants in nature do not experience the same pressures as plants in agriculture. The wide diversity of photosynthetic properties found in nature, including among the wild ancestors of crop plants, in broad terms is expected to represent different optima for photosynthesis adapted to different naturally occurring niches. These niches are, however, often limiting in various ways, for example, in water and nutrients (such as nitrogen and phosphate), and plants in nature usually experience strong competition from their neighbors (Theeuwens et al. 2022). In agriculture, however, competition with noncrop plant species is usually eliminated (or largely so), nutrients and water are often supplied, and attack from pests and diseases are managed, so far as possible, especially in intensive agriculture and protected horticulture. These agricultural environments do not have any perfectly natural analogs, so evolution will not have optimized photosynthesis for agriculture, and it can therefore still be improved in crop plants. In principle, therefore, photosynthesis is no different to those other plant properties that have been improved by domestication and breeding, despite their having been previously optimized by evolution in the wild ancestors of our crop plants. In the past, however, breeding for improved photosynthesis has occurred only to a limited extent, owing to the difficulty of phenotyping for photosynthesis alongside the complexity of the process (Theeuwens et al. 2022).

Variation in photosynthetic properties of land plants is a valuable resource in terms of understanding the operation and limitations of photosynthesis and finding photosynthetic syndromes or traits that could potentially be used to improve crop plant photosynthesis. The contribution that natural variation can make to understanding and improving photosynthesis can be divided into the following five broad categories:

1. The limits to the adaptability of plant photosynthesis to extreme habitats and environments.
2. Trait variation within or across species that facilitates the physiological analysis of traits and the identification of their underlying mechanisms.
3. The co-occurrence of subtraits that reveal syndromes of evolutionary adaptation of photosynthesis.
4. Physiological models or templates that serve as options for improving photosynthesis.
5. Sources of genetic (or allelic) variation that can be used to breed (including by means of novel plant breeding techniques) for improved photosynthesis.

In addition to providing a way to extend our knowledge of the operation and regulation of photosynthesis and the limits reached by evolutionary refinement of the process, the occurrence of natural variation of photosynthesis implies there must be an underlying



genetic underpinning. If we can unravel this genetic basis of variation for photosynthetic traits we can begin to systematically breed for improved photosynthetic traits. The importance of linking variation in photosynthesis (i.e. phenotypic variation) to genetics is paramount whether or not novel plant breeding techniques or conventional breeding is used to genetically improve a plant. The problem is that while the genes for the headline components of photosynthesis (e.g. the subunits of PSII or PSI) are well known, the genetic basis of variation in many key photosynthetic properties is poorly understood, even when the physiology of that variation is well known. For example, the maximum rate of photosynthesis is correlated with numerous changes in the protein, lipid, and cofactor composition of the leaf, alongside anatomical differences (e.g. Osmond et al. 1980; Schulze and Chapin 1987; van Bel and Gamalei 1992; Brodribb et al. 2007; Taylor et al. 2022). Higher rates of photosynthesis are, for example, correlated with greater activities of Cyt  $b_6/f$  complex or Rubisco, but this increase in activity is largely achieved by there being more of these components per unit area and not by there being super Cyt  $b_6/f$  or Rubisco (i.e. complexes with substantially higher specific activity; e.g. Osmond et al. 1980; Makino et al. 1997; Schöttler et al. 2015; Miller et al. 2017). The genetic basis for how an increased amount of these complexes per unit area of leaf is achieved is poorly understood despite phenotypic variation in  $P_{max}$  being widely studied and encountered. The same can be said of other photosynthetic traits (or subtraits). This does not mean, however, that no natural variation exists for genes coding for the headline components of photosynthesis—such variation does exist (e.g. Prins et al. 2016; Prinzenberg et al. 2020) and can potentially be exploited.

If there is variation for a trait within a species, or a pool of species, that can be hybridized to give rise to genetically segregating offspring, then it is possible to correlate phenotypic variation with genomic variation using a mapping population and by this means identify genomic regions—QTL—within which a gene or genes that are causal for phenotypic variation are located (e.g. Harbinson et al. 2012; Theeuwens et al. 2022). Mapping populations can be constructed in various ways (Theeuwens et al. 2022), but critically they need to contain variation for the trait under investigation. The individuals comprising the population (typically 100 and often many more, depending on the nature of the population) must also be genomically mapped using markers (commonly SNPs) that essentially serve to describe the variation of the genome. If there is both genomic and phenotypic variation, these can be correlated and regions of genomic variation that are associated with phenotypic variation identified. A critical requirement is that of phenotyping; photosynthesis is strongly affected by the environment and is, in any case, difficult to measure rapidly and on the large scale needed to adequately phenotype a large population. This can be achieved by the large-scale use of portable gas analyzers or by using robotic chlorophyll fluorescence-based imaging systems (e.g. Flood et al. 2016). Comparing marker-assisted breeding and genomic selection on the one hand with genetic modification and gene editing on the other hand, the latter requires the identification of genes whose variation results in phenotypic difference, while the former can be carried out knowing only the association between phenotypic variation and variation in genomic markers (e.g. single-nucleotide markers or SNPs)—an approach that is widely used in commercial plant breeding [see Theeuwens et al. (2022) for a summary of QTLs and their use].

Improving photosynthesis is still a largely unexplored strategy for enhancing crop productivity, offering a means to expand crop improvement possibilities while sustaining yields. Even considering only the land plants (the embryophytes), photosynthesis has a wide range of properties many of which vary not only from species

to species, but within a species. In some cases, these properties and their variation are not well explored. This natural variation for photosynthesis is, nonetheless, a valuable resource for improving crop photosynthetic properties, providing us with physiological templates and genetic resources with which we can improve crop photosynthesis using either conventional or novel plant breeding techniques. To make best use of this resource, we need to better understand the limitations on photosynthesis in the field and improve the tools needed for identifying phenotypic variation for photosynthesis. We also need better tools for identifying the causal genes underlying photosynthetic variation and, finally, develop strategies for applying these discoveries in crop improvement programs.

## Modeling photosynthesis


### By Xin-Guang Zhu

Engineering canopy photosynthesis, instead of leaf photosynthesis, is required to improve crop yield potential. Canopy photosynthesis is an integral of photosynthetic  $CO_2$  uptake rates for all leaves in a canopy, including those at the top, which usually receive high irradiance, and those in the lower layers, which usually receive low irradiance (Zhu et al. 2012). Earlier wheat breeding programs suggest that cultivars with a higher light-saturated rate of leaf photosynthesis are usually associated with a lower leaf area index, which consequently negates the positive impact of enhancing leaf photosynthetic rate (Evans and Dunstone 1970). Therefore, identifying engineering targets that can increase canopy photosynthesis becomes crucial to achieving the desired goal of increasing photosynthesis for greater yields.

Canopy photosynthesis is controlled by both the microclimate parameters inside a canopy, such as  $CO_2$ , light, temperature, and humidity, and the photosynthetic properties of all leaves in a canopy. Although canopy photosynthesis can be measured with canopy chambers (Song et al. 2016), the development of accurate canopy photosynthesis models is needed for the precise dissection of the main players and regulatory factors. Various models have been developed, often based on 3D structural modeling and energy balance approaches and which differ in their levels of detail in canopy architecture and microclimate heterogeneity (dePury and Farquhar 1997; Song et al. 2017; Liu et al. 2021). These models have been used to identify key architectural parameters, such as optimal leaf area index and ideal leaf angle to enhance canopy photosynthesis (Song et al. 2013; Liu et al. 2021).

Besides microclimate, another factor controlling canopy photosynthesis is leaf photosynthetic rate, which shows large variation among species and between cultivars of the same species (van Bezouw et al. 2019). This variation is mainly attributed to differences in photosynthetic properties, e.g. Rubisco content, Rubisco activation state, CBB cycle enzyme activities, the abundance of electron transfer chain components, mesophyll and stomatal conductance (Long et al. 2015), and leaf anatomical features (Giuliani et al. 2013). To enable the precise dissection of factors controlling leaf photosynthetic rate, 3D reaction–diffusion models of leaf photosynthesis have been developed that effectively couple leaf anatomical features,  $CO_2$  diffusion processes, and light distribution inside a leaf (Xiao et al. 2016, 2022; Xiao and Zhu 2017). These models indicate that leaf biophysical and biochemical properties associated with photosynthesis play a dominant role in determining photosynthetic rates, while leaf anatomy makes a relatively minor contribution (Xiao et al. 2022).

There is a long history of modeling photosynthetic systems to identify critical proteins and enzymes and their biochemical and



**Figure 10.** Systems approach to identify options to engineer photosynthesis for higher efficiency. **A)** Multi-scale models of photosynthesis. Models for photosynthesis at different organismal scales spanning from organelle, cell, leaf, up to canopy scales, have been developed. These models are used to define the architectural, anatomical, biophysical, and biochemical parameters controlling photosynthetic efficiency. **B)** Options to engineering photosynthesis for greater efficiency, divided into three categories: (i) increase the delivery of  $\text{CO}_2$ , (ii) optimize light distribution across a canopy, and (iii) manipulate photosynthetic machinery.

biophysical properties that control photosynthetic efficiency. Several modeling approaches have been used, e.g. systems of ordinary differential equations to model photosynthesis-related metabolic processes (Zhu et al. 2007, 2013), reaction-diffusion models to simulate the gas diffusion and the coupled reaction processes in a MC or leaf (Tholen and Zhu 2011; Tholen et al. 2012; Xiao and Zhu 2017), and ray-tracing algorithms to simulate light distribution inside a leaf (Xiao et al. 2016) (Fig. 10). Systems models of photosynthesis for  $\text{C}_3$ ,  $\text{C}_4$ , and CAM leaves have been developed (Zhu et al. 2013; Wang et al. 2014b, 2023), as well as models simulating stomatal conductance (Buckley et al. 2003) and NPQ dynamics (Zaks et al. 2012).

Analyses with these systems models have generated tremendous insight into strategies for engineering photosynthesis. For example, a dynamic system model of canopy photosynthesis showed that faster recovery from photoprotective states improves canopy photosynthesis (Zhu et al. 2004), which in turn results in increased biomass production in both soybean and tobacco (Kromdijk et al. 2016; De Souza et al. 2022). Systems models of photorespiratory bypass suggest that decreased expression of PLGG1, a glycolate/glycerate transporter (Pick et al. 2013), results in an increased benefit of photorespiratory bypass and further increased photosynthetic  $\text{CO}_2$  uptake rate (Xin et al. 2015), which was later experimentally confirmed in tobacco (South et al. 2019) and rice (Shen et al. 2019a). Analysis with a systems model of  $\text{C}_3$  photosynthesis suggested that simultaneous overexpression of SBPase and FBPase results in increased photosynthesis, which was again later demonstrated in wheat (Zhu et al. 2007; Simkin et al. 2017a). Modeling together with field experiments has established a suite of engineering options that can be used to overcome limitations on photosynthesis. These options are discussed individually in the earlier sections. Here, we summarize them from the angle of access to substrates of photosynthesis, i.e.  $\text{CO}_2$  and light, and limitations due to the inefficiency of proteins or enzymes involved in photosynthesis (Fig. 10B).

Although there are successes of current models in guiding photosynthesis engineering, the suggested strategies do not always deliver the predicted increase in photosynthesis and biomass, as in the case of increasing the NPQ relaxation in *Arabidopsis* (Garcia-Molina and Leister 2020) and potato (Lehretz et al. 2022), or increasing the expression of SBPase in rice (Feng et al. 2007). One potential reason is that factors limiting photosynthesis might shift to different processes under different environments or in

different species. For example, under high light, Rubisco has higher control over photosynthetic  $\text{CO}_2$  uptake, while under low light, components of the electron transfer chain exert higher control (Raines 2022); as a result, engineering a particular enzyme may not necessarily increase photosynthesis for a particular crop under a particular condition. However, this does not nullify the possibility that a particular step limits photosynthetic rates under other conditions, which therefore calls for field studies to test whether a particular engineering option will work for a particular plant or a particular cultivar. In some sense, once the original concept that a particular protein is limiting photosynthesis in model plant species is established, private enterprises might be better positioned to take the lead in systematically testing its application in different crops under different conditions through large-scale field testing.

It is worth pointing out that while the use of the systems model of photosynthesis has shown promise in guiding photosynthesis engineering for higher efficiency, this area of study is in its infancy. Great opportunities lay ahead to develop more advanced models to guide future engineering and design for higher photosynthetic efficiency. First, current models need to include a description of the acclimation of photosynthesis (e.g. expression, assembly, and degradation of proteins) to various environmental factors such as light, humidity, and temperature as well as internal factors like sink capacities. For example, a recent study shows that the ATP required for PSII repair processes is on average 4.6% of that used for photosynthetic carbon assimilation (Yi et al. 2022). Moreover, studies show that increasing the synthesis of the D1 protein, which has a high turnover rate (Li et al. 2018), enhances high photosynthetic rates (Chen et al. 2020). Second, the complete redesign of photosynthetic  $\text{CO}_2$  fixation and carbon metabolism has the potential to dramatically improve photosynthetic efficiency (Bar-Even et al. 2010; Trudeau et al. 2018). Third, systems modeling of excitation energy transfer and electron transfer processes need to incorporate the spatial organization of photosynthetic proteins and pigment-protein complexes in the thylakoid membrane. This information is crucial not only for understanding the principles underlying the high photosynthetic efficiency but also for identifying targets to optimize plants for even higher efficiency. Finally, the enzymes or proteins predicted as limiting factors by systems modeling can be directly correlated with sequence variations and molecular dynamics simulations to identify optimal genomic editing strategies for photosynthesis-related genes. With the rapid increase in computational power and the

expanding capacity of photosynthesis models, we are entering an era of rational design not only for designing new pathways but also new proteins to significantly enhance the efficiency of crops.

## Smart canopy for enhanced crop yield and NUE

By Young B. Cho and Donald R. Ort

In the dense monoculture system of current row crop agriculture, low-light use efficiency at the top of the crop canopy and limited light availability within the canopy conspire to limit carbon gain. The notion of a “smart canopy” concept was proposed for optimizing canopy photosynthesis (Ort et al. 2015). Proposed strategies for improving photosynthesis at the canopy level could benefit from considering a smart canopy concept in the context of optimization of nitrogen (N) distribution within the canopy and to provide promising target genes and enabling technologies.

Studies in optimization theory suggested that maximizing canopy photosynthesis requires N distribution proportional to the light availability within the canopy (Field 1983; Anten et al. 1995). Leaf N decreases gradually from the top to the bottom of the canopy, which is a crucial adaptation to declining light availability within the canopy (Hirose 2005; Pons 2016). The vertical distributions of light and N within canopies are described using the extinction coefficients for light ( $K_L$ ) and N ( $K_N$ ) (Hirose and Werger 1987). Canopy photosynthesis can be maximized by the optimal gradient of N and light (i.e.  $K_N/K_L = 1$ ) within canopies (Anten et al. 1995). A meta-analysis of canopy N distribution has revealed that the  $K_N/K_L$  for most plant species is approximately 0.5 (Hikosaka et al. 2016), implying that improving canopy photosynthesis can be achieved by either reducing  $K_L$  (more uniform light distribution) or increasing  $K_N$  (less uniform N distribution) [see Niinemets (2023) for more details]. More than half of the leaf N is invested in the photosynthetic apparatus (Evans 1989), suggesting that “smart” regulation of photosynthetic apparatus within a canopy will mediate optimizing N distribution.

Various strategies have been proposed to enhance leaf and canopy photosynthesis, with many of them having been tested in field experiments. Evans and Clarke (2019) assessed the N costs of these strategies. For example, engineering Rubisco and ATP synthase was expected to have a high N cost, as they account for a significant portion of the N budget of the leaf (20% and 8%, respectively). On the other hand, overexpressing enzymes such as Cyt  $b_6/f$ , Psbs-VDE-ZEP, SBPase, and FBP aldolase incurred a medium N cost (Evans and Clarke 2019). The concept of a “smart canopy” can minimize the N cost by overexpressing rate-limiting enzymes only at the top of the canopy where sufficient light is available while repressing N investments in the enzymes where light is limited. This smart canopy strategy aims to maximize canopy-level photosynthesis while optimizing N distribution within the canopy. Interestingly, Evans and Clarke (2019) pointed out that one of the strategies for improving photosynthesis can save N, rather than incur additional N costs: reducing the light-capturing machinery such as chlorophyll and antenna proteins. Ort et al. (2015) proposed the smart canopy has more RCs and fewer antenna proteins at the top of the canopy, with the reverse arrangement in the lower canopy.

For agricultural purposes, crop plants overinvest in light capture while underinvesting in light utilization to optimize canopy carbon gain. Studies on low-chlorophyll crops have shown that many plants invest excessively in the production of chlorophyll and its associated LHC (Li et al. 2013; Gu et al. 2017b; Kirst et al. 2017;

Slattery et al. 2017; Sakowska et al. 2018; Cho et al. 2023). One significant evolutionary advantage of overinvesting N in light capture is that it confers a selective advantage by shading potential competitors (Zhang et al. 1999). Even when a leaf is light-saturated and cannot utilize additional light, intercepting more light prevents potential competitors from receiving and benefiting from it; plants even produce more leaves than necessary to capture light (Srinivasan et al. 2017). Another benefit may be that many crops evolved under conditions of limited N, and thus, it is adaptive to sequester N whenever it is available, storing it in proteins like Rubisco to conserve this typically scarce resource (Denison et al. 2003). However, this N investment strategy is suboptimal for densely planted agricultural monocultures (Loomis 1993; Denison et al. 2003), where the main goal is to maximize net primary productivity in the field. A decrease of light capture, rather than an increase, to achieve improved light distribution in the canopy would benefit NUE and perhaps overall canopy photosynthesis (Ort et al. 2011).

Theory suggests an increase in total canopy photosynthesis if N saved by reducing leaf chlorophyll content was optimally reallocated to photosynthetic capacity that matched increased light levels within the reduced chlorophyll canopy (Song et al. 2017; Zhou et al. 2023). Experimental verification is needed to assess the impacts of these modifications on leaf and canopy photosynthesis. Theoretically, a 50% reduction in leaf chlorophyll and LHC could result in 7% to 9% savings in leaf N without negatively affecting canopy photosynthesis (Walker et al. 2018). The saved N from the reduction of chlorophyll could be utilized for the overexpression of rate-limiting enzymes in the CBB cycle, such as Rubisco, FBP aldolase, and SBPase (Zhu et al. 2007). Proposed strategies include different N investment approaches at various canopy levels, allocating N from lower to upper canopy leaves. Beyond the predicted advantages of re-investing saved N in increased photosynthetic capacity, Cho et al. (2023) demonstrated that reducing canopy chlorophyll levels in tobacco by up to 50% did not compromise carbon assimilation but increased seed N concentration by 7%, indicating that saved N from reducing chlorophyll and LHCs may be redirected toward seed N.

Multigene constructs, overexpressing a suite of rate-limiting enzymes such as Rubisco, SBPase, and FBP aldolase at the top of the canopy while downregulating overinvested enzymes in the lower canopy, will better optimize N distribution and improve canopy photosynthesis. These constructs should be regulated by distinct promoters customized for specific canopy heights. Perhaps the ratio of red/far-red (R/FR) light serves as a reasonable inducer, especially as the R/FR ratio decreases with canopy height. With the advent of optogenetics, blue light- and FR light-inducible gene expression systems have been tested in microorganisms (Gligorovski et al. 2023; Liu et al. 2023). However, these systems have not yet been tested in plants. The red light-inducible promoter has been implemented in a plant system, but it was found to turn off gene expression under white light (Ochoa-Fernandez et al. 2020), rendering it unsuitable for the intended purpose. Therefore, a canopy height-specific gene expression system should be further investigated under actual canopy conditions, preferably in field experiments.

The redistribution of the investment of N from light capture to photosynthetic capacity is a legitimate opportunity to improve light use efficiency and canopy carbon gain for the dense canopies of current row crop agriculture. Candidate targets for N savings and reinvestment have been identified and highlighted in this brief overview. The next critical steps involve adapting gene regulation and the validation of these predictions in target crop species in field experiments and ultimately in multi-location field trials to assess impacts on crop yield.

## Outlook

In this perspective, we have discussed key targets and methodologies designed to enhance photosynthetic efficiency, highlighting their significant potential for crop yield improvement. The next critical step is the integration of these approaches. By combining strategies such as optimizing Rubisco efficiency, boosting electron transport, introducing novel pigments and refining canopy structures, and improving stomatal regulation and photosynthetic responses to environmental fluctuations, we can harness their collective potential. Importantly, this integration must take into account the complex interplay of synergistic and antagonistic effects among these modifications to maximize their benefits for agricultural productivity. The combination of these strategies should then aim not only at stacking improvements but also at ensuring real benefits in terms of plant resilience and productivity in diverse farming conditions. This means not just mixing different improvements but also understanding how they work together in field conditions—how they affect plant growth, sink–source relationships, and the response to environmental stresses, and when and how photosynthesis limits crop productivity. This requires extending the focus beyond enhancing photosynthetic efficiency, incorporating traits that increase resilience to abiotic and biotic stresses, improve water use efficiency and NUE, and decrease the yield gap under fluctuating environmental conditions. Importantly, improving primary production creates flexibility in crop plant design. The increased carbon could be used to increase yield or, for example, it could be allocated, without a decrease in yield, to increased root biomass to improve agricultural sustainability by improving nutrient or water capture or by increasing soil organic carbon. The goal is to create crops that are more productive but also better suited to the diverse environments they grow in, directly addressing the pressing demands of global food security.

## Acknowledgments

The authors thank Nan Eckardt for her support in organizing this perspective. The authors apologize to researchers whose important work was not cited due to space limitations. P.P. thanks Paul Hardy for a critical reading of the section on enhancing the efficacy of photosynthesis under field conditions.

## Funding

Work on improving photosynthesis in the R.C. lab was supported by the European Union's Horizon2020 Research and Innovation Programme (grant 862201 CAPITALISE) and the Dutch Organization for Scientific Research (NWO—TOP grant). Work on pale green barley mutants in lab of P.P. was supported by the European Union's Horizon Research and Innovation Actions under grant number 101082091—BEST-CROP. Research on Rieske FeS by M.E. was supported by the ARC Centre of Excellence for Translational Photosynthesis (CE140100015). Work in the laboratories of C.R. and L.T. was supported by the Realising Improved Photosynthetic Efficiency (RIPE) initiative awarded to the University of Essex by University of Illinois, USA. RIPE was possible through support from the Bill & Melinda Gates Foundation, DFID, and FFAR, grant. This work was also supported by the Biotechnology and Biological Sciences Research Council (BBSRC) grants BB/J004138/1, BB/H01960X/1, and BB/N021045/1 and the European Union's Horizon2020 Research and Innovation Programme (no. 862201) project CAPITALISE. T.L. also acknowledge support from Biotechnology and Biological Sciences Research council (BBSRC)

grants; BB/S005080/1; BB/T004274/1; BB/PO27970/1. Work in the E.C.-S. lab was supported by the European Union's Horizon2020 Research and Innovation Programme (grants 862201 CAPITALISE and 862127 PhotoBoost); the project Realizing Increased Photosynthetic Efficiency (RIPE), which is funded by Bill & Melinda Gates Agricultural Innovations grant investment 57248, awarded to Lancaster University by the University of Illinois, USA; and the BBSRC funded BBSRC Institute Strategic Programme: Delivering Sustainable Wheat (DSW; grant BB/X011003/1). Work on engineering pCCMs into plants in lab of A.M. was supported by the BBSRC (grants BB/X018377/1 and BB/S015531/1), the Carbon Technology Research Foundation (grant AP23-1\_023), and a joint award from the Bill & Melinda Gates Foundation and the Foreign, Commonwealth and Development Office (FCDO) awarded to the University of Edinburgh by Princeton University, USA. Research on NPQ by D.P.-T. and K.N. was supported by the US Department of Energy, Office of Science, Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division under Field Work Proposal number 449B, and research on VPZ by D.P.-T. and K.N. was supported by the project Realizing Increased Photosynthetic Efficiency (RIPE), that is funded by Bill & Melinda Gates Agricultural Innovations grant investment 57248, awarded to the University of California, Berkeley by the University of Illinois, USA. D.P.-T. was supported by the Berkeley Fellowship and the National Science Foundation Graduate Research Fellowship Program (grant DGE 1752814). K.K.N. is an investigator of the Howard Hughes Medical Institute. Research in A.W. lab was supported by the European Union's H2020 research and innovation and the Horizon Europe Framework programme (grants GAIN4CROPS, GA No. 862087), and Horizon Europe Framework Programme (Boosting Photosynthesis to deliver novel crops for the circular bioeconomy - BEST-CROP, GA No. 101082091) and the Deutsche Forschungsgemeinschaft [Cluster of Excellence for Plant Sciences (CEPLAS) under Germany's Excellence Strategy EXC-2048/1 under project ID 390686111 and CRC TRR 341 "Plant Ecological Genetics"].

*Conflict of interest statement.* K.N. is an inventor on a patent "Transgenic plants with increased photosynthesis efficiency and growth" US20230183731A1, and K.N. and D.P.-T. are inventors on a patent application "Methods of screening for plant gain of function mutations and compositions therefor" US20230323480A1. All other authors declare no competing interests.

## References

- Adler L, Díaz-Ramos A, Mao Y, Pukacz KR, Fei C, McCormick AJ. New horizons for building pyrenoid-based CO<sub>2</sub>-concentrating mechanisms in plants to improve yields. *Plant Physiol.* 2022;190(3):1609–1627. <https://doi.org/10.1093/plphys/kiac373>
- Amaral J, Lobo AKM, Carmo-Silva E. Regulation of Rubisco activity in crops. *New Phytol.* 2024;241(1):35–51. <https://doi.org/10.1111/nph.19369>
- Ang WSL, How JA, How JB, Mueller-Cajar O. The stickers and spacers of Rubiscondensation: assembling the centrepiece of biophysical CO<sub>2</sub>-concentrating mechanisms. *J Exp Bot.* 2023;74(2):612–626. <https://doi.org/10.1093/jxb/erac321>
- Anten NPR, Schieving F, Werger MJA. Patterns of light and nitrogen distribution in relation to whole canopy carbon gain in C<sub>3</sub> and C<sub>4</sub> mono- and dicotyledonous species. *Oecologia.* 1995;101(4):504–513. <https://doi.org/10.1007/BF00329431>
- Atkinson N, Mao Y, Chan KX, McCormick AJ. Condensation of Rubisco into a proto-pyrenoid in higher plant chloroplasts. *Nat Commun.* 2020;11(1):6303. <https://doi.org/10.1038/s41467-020-20132-0>

- Badger MR, Sharwood RE. Rubisco, the imperfect winner: it's all about the base. *J Exp Bot.* 2023;74(2):562–580. <https://doi.org/10.1093/jxb/erac458>
- Bar-Even A, Noor E, Lewis NE, Milo R. Design and analysis of synthetic carbon fixation pathways. *Proc Natl Acad Sci U S A.* 2010;107(19):8889–8894. <https://doi.org/10.1073/pnas.0907176107>
- Barrett J, Girr P, Mackinder LCM. Pyrenoids: CO<sub>2</sub>-fixing phase separated liquid organelles. *Biochim Biophys Acta Mol Cell Res.* 2021;1868(5):118949. <https://doi.org/10.1016/j.bbamcr.2021.118949>
- Bassi R, Dall'Osto L. Dissipation of light energy absorbed in excess: the molecular mechanisms. *Ann Rev Plant Biol.* 2021;72(1):47–76. <https://doi.org/10.1146/annurev-arplant-071720-015522>
- Bauwe H. Photorespiration—Rubisco's repair crew. *J Plant Physiol.* 2023;280:153899. <https://doi.org/10.1016/j.jplph.2022.153899>
- Bauwe H, Keerberg O, Bassuner R, Parnik T, Bassuner B. Reassimilation of carbon-dioxide by *Flaveria* (Asteraceae) species representing different types of photosynthesis. *Planta.* 1987;172(2):214–218. <https://doi.org/10.1007/BF00394590>
- Beckmann J, Lehr F, Finazzi G, Hankamer B, Posten C, Wobbe L, Kruse O. Improvement of light to biomass conversion by de-regulation of light-harvesting protein translation in *Chlamydomonas reinhardtii*. *J Biotechnol.* 2009;142(1):70–77. <https://doi.org/10.1016/j.jbiotec.2009.02.015>
- Bellasio C, Ermakova M. Reduction of bundle sheath size boosts cyclic electron flow in *C<sub>4</sub> Setaria viridis* acclimated to low light. *Plant J.* 2022;111(5):1223–1237. <https://doi.org/10.1111/tpj.15915>
- Bellasio C, Farquhar GD. A leaf-level biochemical model simulating the introduction of *C<sub>2</sub>* and *C<sub>4</sub>* photosynthesis in *C<sub>3</sub>* rice: gains, losses and metabolite fluxes. *New Phytol.* 2019;223(1):150–166. <https://doi.org/10.1111/nph.15787>
- Benes B, Guan K, Lang M, Long SP, Lynch JP, Marshall-Colon A, Peng B, Schnable J, Sweetlove LJ, Turk MJ. Multiscale computational models can guide experimentation and targeted measurements for crop improvement. *Plant J.* 2020;103(1):21–31. <https://doi.org/10.1111/tpj.14722>
- Bertolino LT, Caine RS, Gray JE. Impact of stomatal density and morphology on water-use efficiency in a changing world. *Front Plant Sci.* 2019;10:225. <https://doi.org/10.3389/fpls.2019.00225>
- Bhat JY, Miličić G, Thieulin-Pardo G, Bracher A, Maxwell A, Ciniawsky S, Mueller-Cajar O, Engen JR, Hartl FU, Wendler P, et al. Mechanism of enzyme repair by the AAA<sup>+</sup> chaperone Rubisco activase. *Mol Cell.* 2017a;67(5):744–756. <https://doi.org/10.1016/j.molcel.2017.07.004>
- Bhat JY, Thieulin-Pardo G, Hartl FU, Hayer-Hartl M. Rubisco activases: AAA<sup>+</sup> chaperones adapted to enzyme repair. *Front Mol Biosci.* 2017b;4:20. <https://doi.org/10.3389/fmolb.2017.00020>
- Björkman O, Holmgren P. Adaptability of the photosynthetic apparatus to light intensity in ecotypes from exposed and shaded habitats. *Physiol Plant.* 1963;16(4):889–914. <https://doi.org/10.1111/j.1399-3054.1963.tb08366.x>
- Blankenship RE, Chen M. Spectral expansion and antenna reduction can enhance photosynthesis for energy production. *Curr Opin Chem Biol.* 2013;17(3):457–461. <https://doi.org/10.1016/j.cbpa.2013.03.031>
- Blikstad C, Dugan EJ, Laughlin TG, Turnšek JB, Liu MD, Shoemaker SR, Vogiatzi N, Remis JP, Savage DF. Identification of a carbonic anhydrase–Rubisco complex within the alpha-carboxysome. *Proc Natl Acad Sci U S A.* 2023;120(43):e2308600120. <https://doi.org/10.1073/pnas.2308600120>
- Borden J, Savage D. New discoveries expand possibilities for carboxysome engineering. *Curr Opin Microbiol.* 2022;61:58–66. <https://doi.org/10.1016/j.mib.2021.03.002>
- Bouvier JW, Emms DM, Kelly S. Rubisco is evolving for improved catalytic efficiency and CO<sub>2</sub> assimilation in plants. *Proc Natl Acad Sci U S A.* 2024;121(11):e2321050121. <https://doi.org/10.1073/pnas.2321050121>
- Bracher A, Whitney SM, Hartl FU, Hayer-Hartl M. Biogenesis and metabolic maintenance of Rubisco. *Ann Rev Plant Biol.* 2017;68(1):29–60. <https://doi.org/10.1146/annurev-arplant-043015-111633>
- Bräutigam A, Schlüter U, Lundgren M, Flachbart S, Ebenhöf O, Schönknecht G, Christin P, Bleuler S, Droz J, Osborne C, et al. Biochemical mechanisms driving rapid fluxes in *C<sub>4</sub>* photosynthesis. *bioRxiv* 387431. <https://doi.org/10.1101/387431>, 9 August 2018, preprint: not peer reviewed
- Brodribb TJ, Feild TS, Jordan GJ. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiol.* 2007;144(4):1890–1898. <https://doi.org/10.1104/pp.107.101352>
- Broncano LS, Pukacz KR, Reichel-Deland V, Schlüter U, Triesch S, Weber APM. Photorespiration is the solution, not the problem. *J Plant Physiol.* 2023;282:153928. <https://doi.org/10.1016/j.jplph.2023.153928>
- Brückner K, Schäfer P, Weber E, Grützner R, Marillonnet S, Tissier A. A library of synthetic transcription activator-like effector-activated promoters for coordinated orthogonal gene expression in plants. *Plant J.* 2015;82(4):707–716. <https://doi.org/10.1111/tpj.12843>
- Buckley CR, Caine RS, Gray JE. Pores for thought: can genetic manipulation of stomatal density protect future rice yields? *Front Plant Sci.* 2020;10:1783. <https://doi.org/10.3389/fpls.2019.01783>
- Buckley TN, Mott KA, Farquhar GD. A hydromechanical and biochemical model of stomatal conductance. *Plant Cell Environ.* 2003;26(10):1767–1785. <https://doi.org/10.1046/j.1365-3040.2003.01094.x>
- Cai BB, Ning Y, Li Q, Li QY, Ai XZ. Effects of the chloroplast fructose-1,6-bisphosphate aldolase gene on growth and low-temperature tolerance of tomato. *Int J Mol Sci.* 2022;23(2):728–742. <https://doi.org/10.3390/ijms23020728>
- Caine RS, Yin X, Sloan J, Harrison EL, Mohammed U, Fulton T, Biswal AK, Dionora J, Chater CC, Coe RA, et al. Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. *New Phytol.* 2019;221(1):371–384. <https://doi.org/10.1111/nph.15344>
- Capó-Bauçà S, Iñiguez C, Aguiló-Nicolau P, Galmés J. Correlative adaptation between Rubisco and CO<sub>2</sub>-concentrating mechanisms in seagrasses. *Nat Plants.* 2022;8(6):706–716. <https://doi.org/10.1038/s41477-022-01171-5>
- Carmo-Silva AE, Salvucci ME. The regulatory properties of Rubisco activase differ among species and affect photosynthetic induction during light transitions. *Plant Physiol.* 2013;161(4):1645–1655. <https://doi.org/10.1104/pp.112.213348>
- Carmo-Silva E, Scales JC, Madgwick PJ, Parry MA. Optimizing Rubisco and its regulation for greater resource use efficiency. *Plant Cell Environ.* 2015;38(9):1817–1832. <https://doi.org/10.1111/pce.12425>
- Chen J-H, Chen S-T, He N-Y, Wang Q-L, Zhao Y, Gao W, Guo F-Q. Nuclear-encoded synthesis of the D1 subunit of photosystem II increases photosynthetic efficiency and crop yield. *Nat Plants.* 2020;6(5):570–580. <https://doi.org/10.1038/s41477-020-0629-z>
- Chen T, Hojka M, Davey P, Sun Y, Dykes GF, Zhou F, Lawson T, Nixon PJ, Lin Y, Liu L-N. Engineering alpha-carboxysomes into plant chloroplasts to support autotrophic photosynthesis. *Nat Commun.* 2023;14(1):2118. <https://doi.org/10.1038/s41467-023-37490-0>
- Chen M, Schliep M, Willows RD, Cai ZL, Neilan BA, Scheer H. A red-shifted chlorophyll. *Science.* 2010;329(5997):1318–1319. <https://doi.org/10.1126/science.1191127>
- Cho YB, Boyd RA, Ren Y, Lee M-S, Jones SI, Ruiz-Vera UM, McGrath JM, Masters MD, Ort DR. Reducing chlorophyll level in seed filling stages results in higher seed nitrogen without impacting canopy

- carbon assimilation. *Plant Cell Environ.* 2023;47(1):1–16. <https://doi.org/10.1111/pce.14737>
- Clapero V, Stitt M, Arrivault S. Natural variation in the metabolism of the Calvin–Benson cycle. *Semin Cell Dev Biol.* 2024;155(Pt A):23–36. <https://doi.org/10.1016/j.semcdb.2023.02.015>
- Clarke JT, Warnock RCM, Donoghue PCJ. Establishing a time-scale for plant evolution. *New Phytol.* 2011;192(1):266–301. <https://doi.org/10.1111/j.1469-8137.2011.03794.x>
- Cowling SB, Treeintong P, Ferguson J, Soltani H, Swarup R, Mayes S, Murchie EH. Out of Africa: characterizing the natural variation in dynamic photosynthetic traits in a diverse population of African rice (*Oryza glaberrima*). *J Exp Bot.* 2022;73(10):3283–3298. <https://doi.org/10.1093/jxb/erab459>
- Croce R, van Amerongen H. Light-harvesting in photosystem I. *Photosynth Res.* 2013;116(2–3):153–166. <https://doi.org/10.1007/s11120-013-9838-x>
- Croce R, van Amerongen H. Light harvesting in oxygenic photosynthesis: structural biology meets spectroscopy. *Science.* 2020;369(6506):eaay2058. <https://doi.org/10.1126/science.aay2058>
- Cutolo EA, Guardini Z, Dall’Osto L, Bassi R. A paler shade of green: engineering cellular chlorophyll content to enhance photosynthesis in crowded environments. *New Phytol.* 2023;239(5):1567–1583. <https://doi.org/10.1111/nph.19064>
- Danila F, Schreiber T, Ermakova M, Hua L, Vlad D, Lo S-F, Chen Y-S, Lambret-Frotte J, Hermanns AS, Athmer B, et al. A single promoter-TALE system for tissue-specific and tuneable expression of multiple genes in rice. *Plant Biotechnol J.* 2022;20(9):1786–1806. <https://doi.org/10.1111/pbi.13864>
- Denison RF, Kiers ET, West SA. Darwinian agriculture: when can humans find solutions beyond the reach of natural selection? *Q Rev Biol.* 2003;78(2):145–168. <https://doi.org/10.1086/374951>
- dePury DGG, Farquhar GD. Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant Cell Environ.* 1997;20(5):537–557. <https://doi.org/10.1111/j.1365-3040.1997.00094.x>
- De Souza AP, Burgess SJ, Doran L, Hansen J, Manukyan L, Maryn N, Gotarkar D, Leonelli L, Niyogi KK, Long SP. Soybean photosynthesis and crop yield are improved by accelerating recovery from photoprotection. *Science.* 2022;377(6608):851–854. <https://doi.org/10.1126/science.adc9831>
- Ding F, Wang M, Zhang S. Overexpression of a Calvin cycle enzyme SBPase improves tolerance to chilling-induced oxidative stress in tomato plants. *Sci Hortic.* 2017;214:27–33. <https://doi.org/10.1016/j.scienta.2016.11.010>
- Dow GJ, Berry JA, Bergmann DC. The physiological importance of developmental mechanisms that enforce proper stomatal spacing in a *Rabidopsis thaliana*. *New Phytol.* 2014;201(4):1205–1217. <https://doi.org/10.1111/nph.12586>
- Drake PL, Froend RH, Franks PJ. Smaller, faster stomata: scaling of stomatal size, rate of response, and stomatal conductance. *J Exp Bot.* 2013;64(2):495–505. <https://doi.org/10.1093/jxb/ers347>
- Drewry DT, Kumar P, Long SP. Simultaneous improvement in productivity, water use, and albedo through crop structural modification. *Glob Chang Biol.* 2014;20(6):1955–1967. <https://doi.org/10.1111/gcb.12567>
- Driever SM, Simkin AJ, Alotaibi S, Fisk SJ, Madgwick PJ, Sparks CA, Jones HD, Lawson T, Parry MAJ, Raines CA. Increased SBPase activity improves photosynthesis and grain yield in wheat grown in greenhouse conditions. *Philos Trans R Soc Lond B Biol Sci.* 2017;372(1730):20160384. <https://doi.org/10.1098/rstb.2016.0384>
- Dunn J, Hunt L, Afsharinafar M, Meselmani MA, Mitchell A, Howells R, Wallington E, Fleming AJ, Gray JE. Reduced stomatal density in bread wheat leads to increased water-use efficiency. *J Exp Bot.* 2019;70(18):4737–4748. <https://doi.org/10.1093/jxb/erz248>
- Eisenhut M, Roell M, Weber APM. Mechanistic understanding of photorespiration paves the way to a new green revolution. *New Phytol.* 2019;223(4):1762–1769. <https://doi.org/10.1111/nph.15872>
- Elias E, Brache K, Schäfers J, Croce R. Coloring outside the lines: exploiting pigment-protein synergy for far-red absorption in plant light-harvesting complexes. *J Am Chem Soc.* 2024a;146(5):3508–3520. <https://doi.org/10.1021/jacs.3c13373>
- Elias E, Liguori N, Saga Y, Schäfers J, Croce R. Harvesting far-red light with plant antenna complexes incorporating chlorophyll d. *Biomacromolecules.* 2021;22(8):3313–3322. <https://doi.org/10.1021/acs.biomac.1c00435>
- Elias E, Oliver TJ, Croce R. Oxygenic photosynthesis in far-red light: strategies and mechanisms. *Annu Rev Phys Chem.* 2024b;75. <https://doi.org/10.1146/annurev-physchem-090722-125847>
- Erb TJ, Zarzycki J. Biochemical and synthetic biology approaches to improve photosynthetic CO<sub>2</sub>-fixation. *Curr Opin Chem Biol.* 2016;34:72–79. <https://doi.org/10.1016/j.cbpa.2016.06.026>
- Erb TJ, Zarzycki J. A short history of RubisCO: the rise and fall of Nature’s predominant CO<sub>2</sub> fixing enzyme. *Curr Opin Biotechnol.* 2018;49:100–107. <https://doi.org/10.1016/j.copbio.2017.07.017>
- Ermakova M, Arrivault S, Giuliani R, Danila F, Alonso-Cantabrana H, Vlad D, Ishihara H, Feil R, Guenther M, Borghi GL, et al. Installation of C<sub>4</sub> photosynthetic pathway enzymes in rice using a single construct. *Plant Biotechnol J.* 2021a;19(3):575–588. <https://doi.org/10.1111/pbi.13487>
- Ermakova M, Bellasio C, Fitzpatrick D, Furbank R, Mamedov F, Von Caemmerer S. Upregulation of bundle sheath electron transport capacity under limiting light in C<sub>4</sub> *Setaria viridis*. *Plant J.* 2021b;106(5):1443–1454. <https://doi.org/10.1111/tjp.15247>
- Ermakova M, Danila FR, Furbank RT, Caemmerer S. On the road to C<sub>4</sub> rice: advances and perspectives. *Plant J.* 2020;101(4):940–950. <https://doi.org/10.1111/tjp.14562>
- Ermakova M, Lopez-Calcagno PE, Raines CA, Furbank RT, Von Caemmerer S. Overexpression of the Rieske FeS protein of the Cytochrome b6f complex increases C<sub>4</sub> photosynthesis in *Setaria viridis*. *Commun Biol.* 2019;2(1):314. <https://doi.org/10.1038/s42003-019-0561-9>
- Ermakova M, Woodford R, Taylor Z, Furbank RT, Belide S, Von Caemmerer S. Faster induction of photosynthesis increases biomass and grain yield in glasshouse-grown transgenic *Sorghum bicolor* overexpressing Rieske FeS. *Plant Biotechnol J.* 2023;21(6):1206–1216. <https://doi.org/10.1111/pbi.14030>
- Evans JR. Photosynthesis and nitrogen relationships in leaves of C<sub>3</sub> plants. *Oecologia.* 1989;78(1):9–19. <https://doi.org/10.1007/BF00377192>
- Evans JR, Clarke VC. The nitrogen cost of photosynthesis. *J Exp Bot.* 2019;70(1):7–15. <https://doi.org/10.1093/jxb/ery366>
- Evans LT, Dunstone RL. Some physiological aspects of evolution in wheat. *Aust J Biol Sci.* 1970;23(4):725–741. <https://doi.org/10.1071/B19700725>
- Faralli M, Lawson T. Natural genetic variation in photosynthesis: an untapped resource to increase crop yield potential. *Plant J.* 2020;101(3):518–528. <https://doi.org/10.1111/tjp.14568>
- Farquhar GD, von Caemmerer S, Berry JA. A biochemical-model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta.* 1980;149(1):78–90. <https://doi.org/10.1007/BF00386231>
- Fei C, Wilson AT, Mangan NM, Wingreen NS, Jonikas MC. Modelling the pyrenoid-based CO<sub>2</sub>-concentrating mechanism provides insights into its operating principles and a roadmap for its engineering into crops. *Nat Plants.* 2022;8(5):583–595. <https://doi.org/10.1038/s41477-022-01153-7>

- Feng L, Wang K, Li Y, Tan Y, Kong J, Li H, Zhu Y. Overexpression of SBPase enhances photosynthesis against high temperature stress in transgenic rice plants. *Plant Cell Rep.* 2007;26(9):1635–1646. <https://doi.org/10.1007/s00299-006-0299-y>
- Ferguson JN, Caproni L, Walter J, Shaw K, Thein MS, Mager S, Taylor G, Cackett L, Mathan J, Vath RL, et al. The genetic basis of dynamic non-photochemical quenching and photosystem II efficiency in fluctuating light reveals novel molecular targets for maize (*Zea mays*) improvement. *bioRxiv*: 565118. <https://doi.org/10.1101/2023.11.01.565118>, 2 November 2023, preprint: not peer reviewed
- Field C. Allocating leaf nitrogen for the maximization of carbon gain: leaf age as a control on the allocation program. *Oecologia.* 1983;56(2–3):341–347. <https://doi.org/10.1007/BF00379710>
- Flecken M, Wang H, Popilka L, Hartl FU, Bracher A, Hayer-Hartl M. Dual functions of a Rubisco activase in metabolic repair and recruitment to carboxysomes. *Cell.* 2020;183(2):457–473. <https://doi.org/10.1016/j.cell.2020.09.010>
- Flood PJ, Harbinson J, Aarts MGM. Natural genetic variation in plant photosynthesis. *Trends Plant Sci.* 2011;16(6):327–335. <https://doi.org/10.1016/j.tplants.2011.02.005>
- Flood PJ, Kruijjer W, Schnabel SK, van der Schoor R, Jalink H, Snel JFH, Harbinson J, Aarts MGM. Phenomics for photosynthesis, growth and reflectance in *Arabidopsis thaliana* reveals circadian and long-term fluctuations in heritability. *Plant Methods.* 2016;12(1):14. <https://doi.org/10.1186/s13007-016-0113-y>
- Förster B, Rourke LM, Weerasooriya HN, Pabuayan ICM, Rolland V, Au EK, Bala S, Bajsa-Hirschel J, Kaines S, Kasili RW. The *Chlamydomonas reinhardtii* chloroplast envelope protein LCIA transports bicarbonate in planta. *J Exp Bot.* 2023;74(12):3651–3666. <https://doi.org/10.1093/jxb/erad116>
- Franks PJ, Farquhar GD. The mechanical diversity of stomata and its significance in gas-exchange control. *Plant Physiol.* 2007;143(1):78–87. <https://doi.org/10.1104/pp.106.089367>
- Freschet GT, Dias ATC, Ackerly DD, Aerts R, van Bodegom PM, Cornwell WK, Dong M, Kurokawa H, Liu G, Onipchenko VG, et al. Global to community scale differences in the prevalence of convergent over divergent leaf trait distributions in plant assemblages. *Glob Ecol Biogeogr.* 2011;20(5):755–765. <https://doi.org/10.1111/j.1466-8238.2011.00651.x>
- Furbank R, Kelly S, von Caemmerer S. Photosynthesis and food security: the evolving story of  $C_4$  rice. *Photosynth Res.* 2023;158(2):121–130. <https://doi.org/10.1007/s11120-023-01014-0>
- Galkin E, Dalal A, Evenko A, Fridman E, Kan I, Wallach R, Moshelion M. Risk-management strategies and transpiration rates of wild barley in uncertain environments. *Physiol Plant.* 2018;164(4):412–428. <https://doi.org/10.1111/pp.12814>
- Gan F, Zhang S, Rockwell NC, Martin SS, Lagarias JC, Bryant DA. Extensive remodeling of a cyanobacterial photosynthetic apparatus in far-red light. *Science.* 2014;345(6202):1312–1317. <https://doi.org/10.1126/science.1256963>
- Garcia-Molina A, Leister D. Accelerated relaxation of photoprotection impairs biomass accumulation in *Arabidopsis*. *Nat Plants.* 2020;6(1):9–12. <https://doi.org/10.1038/s41477-019-0572-z>
- Geiger DR, Servaites JC. Diurnal regulation of photosynthetic carbon metabolism in  $C_3$  plants. *Ann Rev Plant Physiol.* 1994;45(1):235–256. <https://doi.org/10.1146/annurev.pp.45.060194.001315>
- Genesio L, Bassi R, Miglietta F. Plants with less chlorophyll: a global change perspective. *Glob Chang Biol.* 2021;27(5):959–967. <https://doi.org/10.1111/gcb.15470>
- Genesio L, Bright RM, Alberti G, Peressotti A, Delle Vedove G, Incerti G, Toscano P, Rinaldi M, Muller O, Miglietta F. A chlorophyll-deficient, highly reflective soybean mutant: radiative forcing and yield gaps. *Environ Res Lett.* 2020;15(7):074014. <https://doi.org/10.1088/1748-9326/ab865e>
- Gionfriddo M, Rhodes T, Whitney S. Perspectives on improving crop Rubisco by directed evolution. *Semin Cell Dev Biol.* 2024;155(Pt A):37–47. <https://doi.org/10.1016/j.semcdb.2023.04.003>
- Gisriel CJ, Cardona T, Bryant DA, Brudvig GW. Molecular evolution of far-red light-acclimated photosystem II. *Microorganisms.* 2022;10(7):1270 <https://doi.org/10.3390/microorganisms10071270>
- Gisriel CJ, Huang HL, Reiss KM, Flesher DA, Batista VS, Bryant DA, Brudvig GW, Wang J. Quantitative assessment of chlorophyll types in cryo-EM maps of photosystem I acclimated to far-red light. *BBA Adv.* 2021;1:100019. <https://doi.org/10.1016/j.bbadv.2021.100019>
- Gisriel CJ, Shen G, Flesher DA, Kurashov V, Golbeck JH, Brudvig GW, Amin M, Bryant DA. Structure of a dimeric photosystem II complex from a cyanobacterium acclimated to far-red light. *J Biol Chem.* 2023;299(1):102815. <https://doi.org/10.1016/j.jbc.2022.102815>
- Gisriel C, Shen G, Kurashov V, Ho MY, Zhang S, Williams D, Golbeck JH, Fromme P, Bryant DA. The structure of Photosystem I acclimated to far-red light illuminates an ecologically important acclimation process in photosynthesis. *Sci Adv.* 2020;6(6):eaay6415. <https://doi.org/10.1126/sciadv.aay6415>
- Giuliani R, Koteyeva N, Voznesenskaya E, Evans MA, Cousins AB, Edwards GE. Coordination of leaf photosynthesis, transpiration, and structural traits in rice and wild relatives (Genus *Oryza*). *Plant Physiol.* 2013;162(3):1632–1651. <https://doi.org/10.1104/pp.113.217497>
- Gligorovski V, Sadeghi A, Rahi SJ. Multidimensional characterization of inducible promoters and a highly light-sensitive LOV-transcription factor. *Nat Commun.* 2023;14(1):3810. <https://doi.org/10.1038/s41467-023-38959-8>
- Gray A, Liu L, Facette M. Flanking support: how subsidiary cells contribute to stomatal form and function. *Front Plant Sci.* 2020;11:881. <https://doi.org/10.3389/fpls.2020.00881>
- Gu J, Zhou Z, Li Z, Chen Y, Wang Z, Zhang H. Rice (*Oryza sativa* L.) with reduced chlorophyll content exhibit higher photosynthetic rate and efficiency, improved canopy light distribution, and greater yields than normally pigmented plants. *Field Crops Res.* 2017b;200:58–70. <https://doi.org/10.1016/j.fcr.2016.10.008>
- Gu J, Zhou Z, Li Z, Chen Y, Wang Z, Zhang H, Yang J. Photosynthetic properties and potentials for improvement of photosynthesis in pale green leaf rice under high light conditions. *Front Plant Sci.* 2017a;8:1082. <https://doi.org/10.3389/fpls.2017.01082>
- Hamaguchi T, Kawakami K, Shinzawa-Itoh K, Inoue-Kashino N, Itoh S, Ifuku K, Yamashita E, Maeda K, Yonekura K, Kashino Y. Structure of the far-red light utilizing photosystem I of *Acaryochloris marina*. *Nat Commun.* 2021;12(1):2333. <https://doi.org/10.1038/s41467-021-22502-8>
- Harbinson J, Prinzenberg AE, Kruijjer W, Aarts MGM. High throughput screening with chlorophyll fluorescence imaging and its use in crop improvement. *Curr Opin Biotechnol.* 2012;23(2):221–226. <https://doi.org/10.1016/j.copbio.2011.10.006>
- Harbinson J, Yin X. Modelling the impact of improved photosynthetic properties on crop performance in Europe. *Food Energy Secur.* 2023;12(1):e402. <https://doi.org/10.1002/fes3.402>
- Harrison EL, Arce Cubas L, Gray JE, Hepworth C. The influence of stomatal morphology and distribution on photosynthetic gas exchange. *Plant J.* 2020;101(4):768–779. <https://doi.org/10.1111/tpj.14560>
- Hazra S, Henderson JN, Liles K, Hilton MT, Wachter RM. Regulation of ribulose-1, 5-bisphosphate carboxylase/oxygenase (rubisco) activase: product inhibition, cooperativity, and magnesium activation. *J Biol Chem.* 2015;290(40):24222–24236. <https://doi.org/10.1074/jbc.M115.651745>

- He S, Crans VL, Jonikas MC. The pyrenoid: the eukaryotic CO<sub>2</sub>-concentrating organelle. *Plant Cell*. 2023;35(9):3236–3259. <https://doi.org/10.1093/plcell/koad157>
- Hetherington AM, Woodward FI. The role of stomata in sensing and driving environmental change. *Nature*. 2003;424(6951):901–908. <https://doi.org/10.1038/nature01843>
- Heyno E, Ermakova M, Lopez-Calcagno PE, Woodford R, Brown KL, Matthews JSA, Osmond B, Raines C, Von Caemmerer S. Rieske FeS overexpression in tobacco provides increased abundance and activity of cytochrome b<sub>6</sub>f. *Physiol Plant*. 2022;174(6):e13803. <https://doi.org/10.1111/ppl.13803>
- Hibberd JM, Sheehy JE, Langdale JA. Using C<sub>4</sub> photosynthesis to increase the yield of rice—rationale and feasibility. *Curr Opin Plant Biol*. 2008;11(2):228–231. <https://doi.org/10.1016/j.pbi.2007.11.002>
- Hikosaka K, Anten NP, Borjigidai A, Kamiyama C, Sakai H, Hasegawa T, Oikawa S, Iio A, Watanabe M, Koike T, et al. A meta-analysis of leaf nitrogen distribution within plant canopies. *Ann Bot*. 2016;118(2):239–247. <https://doi.org/10.1093/aob/mcw099>
- Hirose T. Development of the Monsi–Saeki theory on canopy structure and function. *Ann Bot*. 2005;95(3):483–494. <https://doi.org/10.1093/aob/mci047>
- Hirose T, Werger MJA. Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy. *Oecologia*. 1987;72(4):520–526. <https://doi.org/10.1007/BF00378977>
- Ho MY, Shen G, Canniffe DP, Zhao C, Bryant DA. Light-dependent chlorophyll f synthase is a highly divergent paralog of PsbA of photosystem II. *Science*. 2016;353(6302):aaf9178. <https://doi.org/10.1126/science.aaf9178>
- Ho MY, Soulier NT, Canniffe DP, Shen G, Bryant DA. Light regulation of pigment and photosystem biosynthesis in cyanobacteria. *Curr Opin Plant Biol*. 2017;37:24–33. <https://doi.org/10.1016/j.pbi.2017.03.006>
- Horaruang W, Klejchová M, Carroll W, Silva-Alvim FA, Waghmare S, Papanatsiou M, Amtmann A, Hills A, Alvim JC, Blatt MR, et al. Engineering a K<sup>+</sup> channel ‘sensory antenna’ enhances stomatal kinetics, water use efficiency and photosynthesis. *Nat Plants*. 2022;8(11):1262–1274. <https://doi.org/10.1038/s41477-022-01255-2>
- Huang G, Peng S, Li Y. Variation of photosynthesis during plant evolution and domestication: implications for improving crop photosynthesis. *J Exp Bot*. 2022;73(14):4886–4896. <https://doi.org/10.1093/jxb/erac169>
- Hylton CM, Rawsthorne S, Smith AM, Jones DA, Woolhouse HW. Glycine decarboxylase is confined to the bundle-sheath cells of leaves of C<sub>3</sub>–C<sub>4</sub> intermediate species. *Planta*. 1988;175(4):452–459. <https://doi.org/10.1007/BF00393064>
- Jansson C, Wullschlegel SD, Kalluri UC, Tuskan GA. Phytosequestration: carbon biosequestration by plants and the prospects of genetic engineering. *Bioscience*. 2010;60(9):685–696. <https://doi.org/10.1525/bio.2010.60.9.6>
- Johnson JE, Berry JA. The role of cytochrome b<sub>6</sub>f in the control of steady-state photosynthesis: a conceptual and quantitative model. *Photosynth Res*. 2021;148(3):101–136. <https://doi.org/10.1007/s11120-021-00840-4>
- Johnson JE, Field CB, Berry J. The limiting factors and regulatory processes that control the environmental responses of C<sub>3</sub>, C<sub>3</sub>–C<sub>4</sub> intermediate, and C<sub>4</sub> photosynthesis. *Oecologia*. 2021;197(4):841–866. <https://doi.org/10.1007/s00442-021-05062-y>
- Kaiser E, Morales A, Harbinson J. Fluctuating light takes crop photosynthesis on a rollercoaster ride. *Plant Physiol*. 2017;176(2):977–989. <https://doi.org/10.1104/pp.17.01250>
- Kasajima I, Ebana K, Yamamoto T, Takahara K, Yano M, Kawai-Yamada M, Uchimiyama H. Molecular distinction in genetic regulation of nonphotochemical quenching in rice. *Proc Natl Acad Sci U S A*. 2011;108(33):13835–13840. <https://doi.org/10.1073/pnas.1104809108>
- Keerbergh O, Pärnik T, Ivanova H, Bassünner B, Bauwe H. C<sub>2</sub> photosynthesis generates about 3-fold elevated leaf CO<sub>2</sub> levels in the C<sub>3</sub>–C<sub>4</sub> intermediate species *Flaveria pubescens*. *J Exp Bot*. 2014;65(13):3649–3656. <https://doi.org/10.1093/jxb/eru239>
- Khaipho-Burch M, Cooper M, Crossa J, de Leon N, Holland J, Lewis R, McCouch S, Murray SC, Rabbi I, Ronald P, et al. Genetic modification can improve crop yields—but stop overselling it. *Nature*. 2023;621(7979):470–473. <https://doi.org/10.1038/d41586-023-02895-w>
- Kirst H, Gabilly ST, Niyogi KK, Lemaux PG, Melis A. Photosynthetic antenna engineering to improve crop yields. *Planta*. 2017;245(5):1009–1020. <https://doi.org/10.1007/s00425-017-2659-y>
- Klimyuk VI, Persello-Cartieaux F, Havaux M, Contard-David P, Schuenemann D, Meierhoff K, Gouet P, Jones JD, Hoffman NE, Nussaume L. A chromodomain protein encoded by the Arabidopsis CAO gene is a plant-specific component of the chloroplast signal recognition particle pathway that is involved in LHCP targeting. *Plant Cell*. 1999;11(1):87–99. <https://doi.org/10.1105/tpc.11.1.87>
- Knudsen S, Wendt T, Dockter C, Thomsen HC, Rasmussen M, Jørgensen ME, Lu Q, Voss C, Murozuka E, Østerberg JT, et al. FIND-IT: accelerated trait development for a green evolution. *Sci Adv*. 2022;8(34):eabq2266. <https://doi.org/10.1126/sciadv.abq2266>
- Kohler IH, Ruiz-Vera UM, VanLoocke A, Thomey ML, Clemente T, Long SP, Ort DR, Bernacchi CJ. Expression of cyanobacterial FBP/SBPase in soybean prevents yield depression under future climate conditions. *J Exp Bot*. 2017;68(3):715–726. <https://doi.org/10.1093/jxb/erw435>
- Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, Long SP. Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science*. 2016;354(6314):857–862. <https://doi.org/10.1126/science.aai8878>
- Küster L, Lücke R, Brabender C, Bethmann S, Jahns P. The amount of zeaxanthin epoxidase but not the amount of violaxanthin de-epoxidase is a critical determinant of zeaxanthin accumulation in *Arabidopsis thaliana* and *Nicotiana tabacum*. *Plant Cell Physiol*. 2023;64(10):1220–1230. <https://doi.org/10.1093/pccp/pcad091>
- Lafferty DJ, Robison TA, Gunadi A, Gunn LH, Van Eck J, Li F-W. Biolistics-mediated transformation of hornworts and its application to study pyrenoid protein localization. *bioRxiv* 563637. <https://doi.org/10.1101/2023.10.23.563637>, 26 October 2023, preprint: not peer reviewed
- Lawson T, Blatt M. Stomatal size, speed and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiol*. 2014;164(4):1556–1570. <https://doi.org/10.1104/pp.114.237107>
- Lawson T, Kramer DM, Raines CA. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. *Curr Opin Biotechnol*. 2012;23(2):215–220. <https://doi.org/10.1016/j.copbio.2011.12.012>
- Lawson T, Milliken AL. Photosynthesis—beyond the leaf. *New Phytol*. 2022;238(1):55–56. <https://doi.org/10.1111/nph.18671>
- Lawson T, Morison JIL. Stomatal function and physiology. In: **Hemsley AR, Poole I**, editors. *The evolution of plant physiology: from whole plants to ecosystem*. Cambridge: Elsevier Academic Press; 2004. p. 217–242.
- Lawson T, Vialet-Chabrand S. Speedy stomata, photosynthesis and plant water use efficiency. *New Phytol*. 2019;221(1):93–98. <https://doi.org/10.1111/nph.15330>
- Lawson T, vonCaemmerer S, Baroli I. Photosynthesis and stomatal behaviour. *Prog Bot*. 2010;72:265–304. [https://doi.org/10.1007/978-3-642-13145-5\\_11](https://doi.org/10.1007/978-3-642-13145-5_11)



- Leakey ADB, Ferguson JN, Pignon CP, Wu A, Jin Z, Hammer GL, Lobell DB. Water use efficiency as a constraint and target for improving the resilience and productivity of C<sub>3</sub> and C<sub>4</sub> crops. *Annu Rev Plant Biol* 2019;70(1):781–808. <https://doi.org/10.1146/annurev-arplant-042817-040305>
- Leegood RC. C<sub>4</sub> photosynthesis: principles of CO<sub>2</sub> concentration and prospects for its introduction into C<sub>3</sub> plants. *J Exp Bot*. 2002;53(369):581–590. <https://doi.org/10.1093/jexbot/53.369.581>
- Lefebvre S, Lawson T, Fryer M, Zakhleniuk OV, Lloyd JC, Raines CA. Increased sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis and growth from an early stage in development. *Plant Physiol*. 2005;138(1):451–460. <https://doi.org/10.1104/pp.104.055046>
- Lehretz GG, Schneider A, Leister D, Sonnwald U. High non-photochemical quenching of VPZ transgenic potato plants limits CO<sub>2</sub> assimilation under high light conditions and reduces tuber yield under fluctuating light. *J Integr Plant Biol*. 2022;64(9):1821–1832. <https://doi.org/10.1111/jipb.13320>
- Leister D. Enhancing the light reactions of photosynthesis: strategies, controversies, and perspectives. *Mol Plant*. 2023;16(1):4–22. <https://doi.org/10.1016/j.molp.2022.08.005>
- Lemonnier P, Lawson T. Calvin cycle and guard cell metabolism impact stomatal function. *Semin Cell Dev Biol*. 2023;155:59–70.
- Levey M, Timm S, Mettler-Altman T, Borghi GL, Koczor M, Arrivault S, Weber AP, Bauwe H, Gowik U, Westhoff P. Efficient 2-phosphoglycolate degradation is required to maintain carbon assimilation and allocation in the C<sub>4</sub> plant *Flaveria bidentis*. *J Exp Bot*. 2019;70(2):575–587. <https://doi.org/10.1093/jxb/ery370>
- Li L, Aro EM, Millar AH. Mechanisms of photodamage and protein turnover in photoinhibition. *Trends Plant Sci*. 2018;23(8):667–676. <https://doi.org/10.1016/j.tplants.2018.05.004>
- Li Y-Y, Guo L-N, Liang C-Z, Meng Z-G, Tahira S, Guo S-D, Zhang R. Overexpression of *Brassica napus* cytosolic fructose-1,6-bisphosphatase and sedoheptulose-1,7-bisphosphatase genes significantly enhanced tobacco growth and biomass. *J Integr Agric*. 2022;21(1):49–59. [https://doi.org/10.1016/S2095-3119\(20\)63438-4](https://doi.org/10.1016/S2095-3119(20)63438-4)
- Li Y, Ren B, Gao L, Ding L, Jiang D, Xu X, Shen Q, Guo S. Less chlorophyll does not necessarily restrain light capture ability and photosynthesis in a chlorophyll-deficient rice mutant. *J Agron Crop Sci*. 2013;199(1):49–56. <https://doi.org/10.1111/j.1439-037X.2012.00519.x>
- Lin MT, Salihovic H, Clark FK, Hanson MR. Improving the efficiency of Rubisco by resurrecting its ancestors in the family Solanaceae. *Sci Adv*. 2022;8(15):eabm6871. <https://doi.org/10.1126/sciadv.abm6871>
- Liu F, Song Q, Zhao J, Mao L, Bu H, Hu Y, Zhu XG. Canopy occupation volume as an indicator of canopy photosynthetic capacity. *New Phytol*. 2021;232(2):941–956. <https://doi.org/10.1111/nph.17611>
- Long SP, Ainsworth EA, Leakey ADB, Nosberger J, Ort DR. Rising atmospheric carbon dioxide: plants FACE the future. *Annu Rev Plant Biol*. 2004;55:591–628. <https://doi.org/10.1146/annurev.arplant.55.031903.141610>
- Long S, Burgess S, Causton I. Redesigning crop photosynthesis. In: **Zeigler R**, editor. *Sustaining Global Food Security: The Nexus of Science and Policy Book*. Australia: CISCO; 2019. p. 131–141.
- Long SP, Humphries S, Falkowski PG. Photoinhibition of photosynthesis in nature. *Annu Rev Plant Biol*. 1994;45(1):633–662. <https://doi.org/10.1146/annurev.pp.45.060194.003221>
- Long SP, Marshall AM, Zhu XG. Engineering crop photosynthesis and yield potential to meet global food demand of 2050. *Cell*. 2015;161(1):56–66. <https://doi.org/10.1016/j.cell.2015.03.019>
- Long SP, Taylor SH, Burgess SJ, Carmo-Silva E, Lawson T, De Souza AP, Leonelli L, Wang Y. Into the shadows and back into sunlight: photosynthesis in fluctuating light. *Annu Rev Plant Biol*. 2022;73(1):617–648. <https://doi.org/10.1146/annurev-arplant-070221-024745>
- Loomis RS. Optimization theory and crop improvement. *Int Crop Sci*. 1993;1:583–588. <https://doi.org/10.2135/1993.internationalcrop.science.c92>
- Lopez-Calcagno PE, Brown KL, Simkin AJ, Fisk SJ, Vialet-Chabrand S, Lawson T, Raines CA. Stimulating photosynthetic processes increases productivity and water-use efficiency in the field. *Nat Plants*. 2020;6(8):1054. <https://doi.org/10.1038/s41477-020-0740-1>
- Löwe H, Kremling A. In-depth computational analysis of natural and artificial carbon fixation pathways. *Bioes Res*. 2021;2021:9898316. <https://doi.org/10.34133/2021/9898316>
- Lundgren MR. C<sub>2</sub> photosynthesis: a promising route towards crop improvement? *New Phytol*. 2020;228(6):1734–1740. <https://doi.org/10.1111/nph.16494>
- Makino A. Photosynthesis improvement for enhancing productivity in rice. *Soil Sci Plant Nutr*. 2021;67(5):513–519. <https://doi.org/10.1080/00380768.2021.1966290>
- Makino A, Sato T, Nakano H, Mae T. Leaf photosynthesis, plant growth and nitrogen allocation in rice under different irradiances. *Planta*. 1997;203(3):390–398. <https://doi.org/10.1007/s004250050205>
- Mallmann J, Heckmann D, Bräutigam A, Lercher MJ, Weber APM, Westhoff P, Gowik U. The role of photorespiration during the evolution of C<sub>4</sub> photosynthesis in the genus *Flaveria*. *Elife*. 2014;3:e02478. <https://doi.org/10.7554/eLife.02478>
- Malnoë A, Schultink A, Shahrabi S, Rumeau D, Havaux M, Niyogi KK. The plastid lipocalin LCNP is required for sustained photoprotective energy dissipation in Arabidopsis. *Plant Cell*. 2018;30(1):196–208. <https://doi.org/10.1105/tpc.17.00536>
- Malone LA, Proctor MS, Hitchcock A, Hunter CN, Johnson MP. Cytochrome b6f—Orchestrator of photosynthetic electron transfer. *Biochim Biophys Acta*. 2021;1862(5):148380. <https://doi.org/10.1016/j.bbabi.2021.148380>
- Mao Y, Catherall E, Díaz-Ramos A, Greiff GR, Azinas S, Gunn L, McCormick AJ. The small subunit of Rubisco and its potential as an engineering target. *J Exp Bot*. 2023;74(2):543–561. <https://doi.org/10.1093/jxb/erac309>
- Mascoli V, Bersanini L, Croce R. Far-red absorption and light-use efficiency trade-offs in chlorophyll f photosynthesis. *Nat Plants*. 2020;6(8):1044–1053. <https://doi.org/10.1038/s41477-020-0718-z>
- Mascoli V, Bhatti AF, Bersanini L, van Amerongen H, Croce R. The antenna of far-red absorbing cyanobacteria increases both absorption and quantum efficiency of Photosystem II. *Nat Commun*. 2022;13(1):3562. <https://doi.org/10.1038/s41467-022-31099-5>
- Matthew JAS, Lawson T. Climate change and stomatal physiology. *Annu Plant Rev*. 2019;2(3):713–752. <https://doi.org/10.1002/9781119312994.apr0667>
- McAusland L, Vialet-Chabrand S, Davey PA, Baker NR, Brendel O, Lawson T. Effects of kinetics of light light-induced stomatal responses on photosynthesis and water-use efficiency. *New Phytol*. 2016;211(4):1209–1220. <https://doi.org/10.1111/nph.14000>
- Miller MAE, O’Cualain R, Selley J, Knight D, Karim MF, Hubbard SJ, Johnson GN. Dynamic acclimation to high light in Arabidopsis thaliana involves widespread reengineering of the leaf proteome. *Front Plant Sci*. 2017;8:1239. <https://doi.org/10.3389/fpls.2017.01239>
- Mirkovic T, Ostroumov EE, Anna JM, van Grondelle R, Govindjee D, Scholes GD. Light absorption and energy transfer in the antenna complexes of photosynthetic organisms. *Chem Rev*. 2017;117(2):249–293. <https://doi.org/10.1021/acs.chemrev.6b00002>
- Miyagawa Y, Tamoi M, Shigeoka S. Overexpression of a cyanobacterial fructose-1,6-/sedoheptulose-1, 7-bisphosphatase in tobacco enhances photosynthesis and growth. *Nat Biotechnol*. 2001;19(10):965–969. <https://doi.org/10.1038/nbt1001-965>

- Miyashita H, Ikemoto H, Kurano N, Adachi K, Chihara M, Miyachi S. Chlorophyll d as a major pigment. *Nature*. 1996;383(6599):402–402. <https://doi.org/10.1038/383402a0>
- Mooney HA, Ehleringer J, Berry JA. High photosynthetic capacity of a winter annual in death valley. *Science*. 1976;194(4262):322–324. <https://doi.org/10.1126/science.194.4262.322>
- Mueller-Cajar O. The diverse AAA+ machines that repair inhibited Rubisco active sites. *Front Mol Biosci*. 2017;4:31. <https://doi.org/10.3389/fmolb.2017.00031>
- Munekage Y. Light harvesting and chloroplast electron transport in NADP-malic enzyme type C<sub>4</sub> plants. *Curr Opin Plant Biol*. 2016;31:9–15. <https://doi.org/10.1016/j.pbi.2016.03.001>
- Nakajima Y, Ueda R. Improvement of photosynthesis in dense microalgal suspension by reduction of light harvesting pigments. *J Appl Phycol*. 1997;9:503–510. <https://doi.org/10.1023/A:1007920025419>
- Nam O, McKenzie C, Dowle A, Dowson M, Barrett J, Mackinder LCM. A protein blueprint of the diatom CO<sub>2</sub>-fixing organelle. *bioRxiv* 564148. <https://doi.org/10.1101/2023.10.26.564148>, 26 October 2023, preprint: not peer reviewed
- Nguyen ND, Pulsford SB, Long BM. Plant-based carboxysomes: another step toward increased crop yields. *Trends Biochem Sci*. 2023;48(10):832–834. <https://doi.org/10.1016/j.tibs.2023.07.003>
- Niinemets Ü. Variation in leaf photosynthetic capacity within plant canopies: optimization, structural, and physiological constraints and inefficiencies. *Photosynth Res*. 2023;158(2):131–149. <https://doi.org/10.1007/s11220-023-01043-9>
- Nilkens M, Kress E, Lambrev P, Miloslavina Y, Müller M, Holzwarth AR, Jahns P. Identification of a slowly inducible zeaxanthin-dependent component of non-photochemical quenching of chlorophyll fluorescence generated under steady-state conditions in *Arabidopsis*. *Biochim Biophys Acta*. 2010;1797(4):466–475. <https://doi.org/10.1016/j.bbabi.2010.01.001>
- Niyogi KK. PHOTOPROTECTION REVISITED: genetic and molecular approaches. *Annu Rev Plant Physiol Plant Mol Biol*. 1999;50(1):333–359. <https://doi.org/10.1146/annurev.arplant.50.1.333>
- Nobel PS. Achievable productivities of certain CAM plants: basis for high values compared with C<sub>3</sub> and C<sub>4</sub> plants. *New Phytol*. 1991;119(2):183–205. <https://doi.org/10.1111/j.1469-8137.1991.tb01022.x>
- Nürnberg DJ, Morton J, Santabarbara S, Telfer A, Joliot P, Antonaru LA, Ruban AV, Cardona T, Krausz E, Boussac A, et al. Photochemistry beyond the red limit in chlorophyll f-containing photosystems. *Science*. 2018;360(6394):1210–1213. <https://doi.org/10.1126/science.aar8313>
- Ochoa-Fernandez R, Abel NB, Wieland FG, Schlegel J, Koch LA, Miller JB, Engesser R, Giuriani G, Brandl SM, Timmer J, et al. Optogenetic control of gene expression in plants in the presence of ambient white light. *Nat Methods*. 2020;17(7):717–725. <https://doi.org/10.1038/s41592-020-0868-y>
- Oh ZG, Ang WSL, Poh CW, Lai S-K, Sze SK, Li H-Y, Bhushan S, Wunder T, Mueller-Cajar O. A linker protein from a red- type pyrenoid phase separates with Rubisco via oligomerizing sticker motifs. *Proc Natl Acad Sci U S A*. 2023a;120(25):e2304833120. <https://doi.org/10.1073/pnas.2304833120>
- Oh ZG, Askey B, Gunn LH. Red Rubiscos and opportunities for engineering green plants. *J Exp Bot*. 2023b;74(2):520–542. <https://doi.org/10.1093/jxb/erac349>
- Orr DJ, Robijns AK, Baker CR, Niyogi KK, Carmo-Silva E. Dynamics of Rubisco regulation by sugar phosphate derivatives and their phosphatases. *J Exp Bot*. 2023;74(2):581–590. <https://doi.org/10.1093/jxb/erac386>
- Ort DR, Merchant SS, Alric J, Barkan A, Blankenship RE, Bock R, Croce R, Hanson MR, Hibberd JM, Long SP, et al. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proc Natl Acad Sci U S A*. 2015;112(28):8529–8536. <https://doi.org/10.1073/pnas.1424031112>
- Ort DR, Zhu X, Melis A. Optimizing antenna size to maximize photosynthetic efficiency. *Plant Physiol*. 2011;155(1):79–85. <https://doi.org/10.1104/pp.110.165886>
- Osmond CB, Björkman O, Anderson DJ. *Physiological processes in plant ecology: toward a synthesis with Atriplex*. vol. 36. Berlin: Springer; 1980
- Papanatsiou M, Amtmann A, Blatt MR. Stomatal spacing safeguards stomatal dynamics by facilitating guard cell ion transport independent of the epidermal solute reservoir. *Plant Physiol*. 2016;172(1):254–263. <https://doi.org/10.1104/pp.16.00850>
- Papanatsiou M, Petersen J, Henderson L, Wang Y, Christie JM, Blatt MR. Optogenetic manipulation of stomatal kinetics improves carbon assimilation, water use, and growth. *Science*. 2019;363(6434):1456–1459. <https://doi.org/10.1126/science.aaw0046>
- Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM. Rubisco activity and regulation as targets for crop improvement. *J Exp Bot*. 2012;64(3):717–730. <https://doi.org/10.1093/jxb/ers336>
- Patel-Tupper D, Kelikian A, Leipertz A, Maryn N, Tjahjadi M, Karavolias NG, Cho M-J, Niyogi KK. Multiplexed CRISPR/Cas9 mutagenesis of rice PSBS1 non-coding sequences for transgene-free overexpression. *Sci Adv*. 2024;in press.
- Paul MJ. Improving photosynthetic metabolism for crop yields: what is going to work? *Front Plant Sci*. 2021;12:743862. <https://doi.org/10.3389/fpls.2021.743862>
- Perin G, Bellan A, Michelberger T, Lyska D, Wakao S, Niyogi KK, Morosinotto T. Modulation of xanthophyll cycle impacts biomass productivity in the marine microalga *Nannochloropsis*. *Proc Natl Acad Sci U S A*. 2023;120(25):e2214119120. <https://doi.org/10.1073/pnas.2214119120>
- Perrine Z, Negi S, Sayre RT. Optimization of photosynthetic light energy utilization by microalgae. *Algal Res*. 2012;1(2):134–142. <https://doi.org/10.1016/j.algal.2012.07.002>
- Pick TR, Bräutigam A, Schulz MA, Obata T, Fernie AR, Weber APM. PLGG1, a plastidic glycolate glycerate transporter, is required for photorespiration and defines a unique class of metabolite transporters. *Proc Natl Acad Sci U S A*. 2013;110(8):3185–3190. <https://doi.org/10.1073/pnas.1215142110>
- Pons TL. Regulation of leaf traits in canopy gradients. In: **Hikosaka K, Niinemets Ü, Anten NPR**, editors. *Canopy photosynthesis: from basics to applications*. Dordrecht, Heidelberg, New York, London: Springer; 2016. p. 143–168.
- Price GD, Caemmerer SV, Evans JR, Siebke K, Anderson JM, Badger MR. Photosynthesis is strongly reduced by antisense suppression of chloroplastic cytochrome b<sub>f</sub> complex in transgenic tobacco. *Funct Plant Biol*. 1998;25(4):445–452. <https://doi.org/10.1071/PP97164>
- Price GD, Pengelly JLL, Forster B, Du J, Whitney SM, von Caemmerer S, Badger MR, Howitt SM, Evans JR. The cyanobacterial CCM as a source of genes for improving photosynthetic CO<sub>2</sub> fixation in crop species. *J Ex Bot*. 2013;64(3):753–768. <https://doi.org/10.1093/jxb/ers257>
- Price GD, Yu J, Caemmerer SV, Evans J, Chow W, Anderson J, Hurry V, Badger M. Chloroplast cytochrome b<sub>f</sub> and ATP synthase complexes in tobacco: transformation with antisense RNA against nuclear-encoded transcripts for the Rieske FeS and ATPD polypeptides. *Funct Plant Biol*. 1995;22(2):285–297. <https://doi.org/10.1071/PP950285>
- Prins A, Orr DJ, Andralojc PJ, Reynolds MP, Carmo-Silva E, Parry MAJ. Rubisco catalytic properties of wild and domesticated relatives

- provide scope for improving wheat photosynthesis. *J Exp Bot.* 2016;67(6):1827–1838. <https://doi.org/10.1093/jxb/erv574>
- Prinzenberg AE, Campos-Dominguez L, Kruijer W, Harbinson J, Aarts MGM. Natural variation of photosynthetic efficiency in *Arabidopsis thaliana* accessions under low temperature conditions. *Plant Cell Environ.* 2020;43(8):2000–2013. <https://doi.org/10.1111/pce.13811>
- Prywes N, Phillips NR, Oltrogge LM, de Pins B, Cowan AE, Taylor-Kearney LJ, Chang HA, Hall LN, Bhatt A, Shih P, et al. Mapping the biochemical landscape of rubisco. *bioRxiv.* 559826. <https://doi.org/10.1101/2023.09.27.559826>, 27 September 2023, preprint: not peer reviewed
- Prywes N, Phillips NR, Tuck OT, Valentin-Alvarado LE, Savage DF. Rubisco function, evolution, and engineering. *Ann Rev Biochem.* 2023b;92(1):385–410. <https://doi.org/10.1146/annurev-biochem-040320-101244>
- Qu Y, Mueller-Cajar O, Yamori W. Improving plant heat tolerance through modification of Rubisco activase in  $C_3$  plants to secure crop yield and food security in a future warming world. *J Exp Bot.* 2023;74(2):591–599. <https://doi.org/10.1093/jxb/erac340>
- Raines CA. The Calvin cycle revisited. *Photosynth Res.* 2003;75(1):1–10. <https://doi.org/10.1023/A:1022421515027>
- Raines CA. Improving plant productivity by retuning regeneration of RuBP in the Calvin Benson Bassham cycle. *New Phytol.* 2022;236(2):350–356. <https://doi.org/10.1111/nph.18394>
- Raissig MT, Matos JL, Anleu Gil MX, Kornfeld A, Bettadapur A, Abrash E, Allison HR, Badgley G, Vogel JP, Berry JA, et al. Mobile MUTE specifies subsidiary cells to build physiologically improved grass stomata. *Science.* 2017;355(6330):1215–1218. <https://doi.org/10.1126/science.aal3254>
- Rasmussen B, Fletcher IR, Brocks JJ, Kilburn MR. Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature.* 2008;455(7216):1101–1104. <https://doi.org/10.1038/nature07381>
- Rawsthorne S, Hylton CM, Smith AM, Woolhouse HW. Distribution of photorespiratory enzymes between bundle-sheath and mesophyll cells in leaves of the  $C_3$ - $C_4$  intermediate species *Moricandia arvensis* (L.) DC. *Planta.* 1988;176(4):527–532. <https://doi.org/10.1007/BF00397660>
- Rosenthal DM, Locke AM, Khozaei M, Raines CA, Long SP, Ort DR. Over-expressing the  $C_3$  photosynthesis cycle enzyme Sedoheptulose-1-7 Bisphosphatase improves photosynthetic carbon gain and yield under fully open air  $CO_2$  fumigation (FACE). *BMC Plant Biol.* 2011;11(1):123. <https://doi.org/10.1186/1471-2229-11-123>
- Rotasperti L, Sansoni F, Mizzotti C, Tadini L, Pesaresi P. Barley's second spring as a model organism for chloroplast research. *Plants.* 2020;9(7):803. <https://doi.org/10.3390/plants9070803>
- Rotasperti L, Tadini L, Chiara M, Crosatti C, Guerra D, Tagliani A, Forlani S, Ezquer I, Horner DS, Jahns P, et al. The barley mutant *happy under the sun 1 (hus1)*: an additional contribution to pale green crops. *Environ Exp Bot.* 2022;196:104795. <https://doi.org/10.1016/j.envexpbot.2022.104795>
- Rungrat T, Almonte AA, Cheng R, Gollan PJ, Stuart T, Aro E, Borevitz JO, Pogson B, Wilson PB. A genome-wide association study of non-photochemical quenching in response to local seasonal climates in *Arabidopsis thaliana*. *Plant Direct.* 2019;3(5):e00138. <https://doi.org/10.1002/pld3.138>
- Sage RF, Khoshravesh R, Sage TL. From proto-Kranz to  $C_4$  Kranz: building the bridge to  $C_4$  photosynthesis. *J Exp Bot.* 2014;65(13):3341–3356. <https://doi.org/10.1093/jxb/eru180>
- Sage RF, Monson RK, Ehleringer JR, Adachi S, Pearcy RW. Some like it hot: the physiological ecology of  $C_4$  plant evolution. *Oecologia.* 2018;187(4):941–966. <https://doi.org/10.1007/s00442-018-4191-6>
- Sage RF, Sage TL, Kocacinar F. Photorespiration and the evolution of  $C_4$  photosynthesis. *Plant Biol.* 2012;63(1):19–47. <https://doi.org/10.1146/annurev-arplant-042811-105511>
- Sahay S, Grzybowski M, Schnable JC, Głowacka K. Genetic control of photoprotection and photosystem II operating efficiency in plants. *New Phytol.* 2023;239(3):1068–1082. <https://doi.org/10.1111/nph.18980>
- Sakoda K, Yamori W, Shimada T, Sugano SS, Hara-Nishimura I, Tanaka Y. Higher stomatal density improves photosynthetic induction and biomass production in *Arabidopsis* under fluctuating light. *Front Plant Sci.* 2020;11:589603. <https://doi.org/10.3389/fpls.2020.589603>
- Sakowska K, Alberti G, Genesio L, Peressotti A, Delle Vedove G, Gianelle D, Colombo R, Rodeghiero A, Panigada C, Juszcak R, et al. Leaf and canopy photosynthesis of a chlorophyll deficient soybean mutant. *Plant Cell Environ.* 2018;41(6):1427–1437. <https://doi.org/10.1111/pce.13180>
- Savir Y, Noor E, Milo R, Tlustý T. Cross-species analysis traces adaptation of Rubisco toward optimality in a low-dimensional landscape. *Proc Natl Acad Sci U S A.* 2010;107(8):3475–3480. <https://doi.org/10.1073/pnas.0911663107>
- Scafaro AP, Posch BC, Evans JR, Farquhar GD, Atkin OK. Rubisco deactivation and chloroplast electron transport rates co-limit photosynthesis above optimal leaf temperature in terrestrial plants. *Nat Commun.* 2023;14(1):2820. <https://doi.org/10.1038/s41467-023-38496-4>
- Schiller K, Bräutigam A. Engineering of crassulacean acid metabolism. *Annu Rev Plant Biol.* 2021;72(1):77–103. <https://doi.org/10.1146/annurev-arplant-071720-104814>
- Schlüter U, Weber APM. Regulation and evolution of  $C_4$  photosynthesis. *Annu Rev Plant Biol.* 2020;71(1):1–33. <https://doi.org/10.1146/annurev-arplant-042916-040915>
- Schöttler MA, Tóth SZ, Boulouis A, Kahlau S. Photosynthetic complex stoichiometry dynamics in higher plants: biogenesis, function, and turnover of ATP synthase and the cytochrome b6f complex. *J Exp Bot.* 2015;66(9):2373–2400. <https://doi.org/10.1093/jxb/eru495>
- Schulze ED, Chapin FS. Plant specialization to environments of different resource availability. In: **Schulze E-D, Zwölfer H**, editors. *Potentials and limitations of ecosystem analysis*. Berlin: Springer; 1987. p. 120–148.
- Sedelnikova OV, Hughes TE, Langdale JA. Understanding the genetic basis of  $C_4$  Kranz anatomy with a view to engineering  $C_3$  crops. *Annu Rev Genet.* 2018;52(1):249–270. <https://doi.org/10.1146/annurev-genet-120417-031217>
- Sharkey TD. Discovery of the canonical Calvin–Benson cycle. *Photosynth Res.* 2019;140(2):235–252. <https://doi.org/10.1007/s11220-018-0600-2>
- Sharkey TD. The discovery of rubisco. *J Exp Bot.* 2023;74(2):510–519. <https://doi.org/10.1093/jxb/erac254>
- Sharwood RE. Engineering chloroplasts to improve Rubisco catalysis: prospects for translating improvements into food and fiber crops. *New Phytol.* 2017;213(2):494–510. <https://doi.org/10.1111/nph.14351>
- Sheehy JE, Dionora MJA, Mitchell PL. Spikelet numbers, sink size and potential yield in rice. *Field Crop Res.* 2001;71(2):77–85. [https://doi.org/10.1016/S0378-4290\(01\)00145-9](https://doi.org/10.1016/S0378-4290(01)00145-9)
- Shen G, Canniffe DP, Ho MY, Kurashov V, van der Est A, Golbeck JH, Bryant DA. Characterization of chlorophyll f synthase heterologously produced in *Synechococcus* sp. PCC 7002. *Photosynth Res.* 2019b;140(1):77–92. <https://doi.org/10.1007/s11220-018-00610-9>
- Shen BR, Wang LM, Lin XL, Yao Z, Xu HW, Zhu CH, Teng HY, Cui LL, Liu EE, Zhang JJ, et al. Engineering a new chloroplastic

- photorespiratory bypass to increase photosynthetic efficiency and productivity in rice. *Mol Plant*. 2019a;12(2):199–214. <https://doi.org/10.1016/j.molp.2018.11.013>
- Simkin AJ, Lopez-Calcagno PE, Davey PA, Headland LR, Lawson T, Timm S, Bauwe H, Raines CA. Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphate aldolase and the photorespiratory glycine decarboxylase-H protein increases CO<sub>2</sub> assimilation, vegetative biomass and seed yield in Arabidopsis. *Plant Biotechnol J*. 2017a;15(7):805–816. <https://doi.org/10.1111/pbi.12676>
- Simkin AJ, Lopez-Calcagno PE, Raines CA. Feeding the world: improving photosynthetic efficiency for sustainable crop production. *J Exp Bot*. 2019;70(4):1119–1140. <https://doi.org/10.1093/jxb/ery445>
- Simkin AJ, McAusland L, Headland LR, Lawson T, Raines CA. Multigene manipulation of photosynthetic carbon assimilation increases CO<sub>2</sub> fixation and biomass yield in tobacco. *J Exp Bot*. 2015;66(13):4075–4090. <https://doi.org/10.1093/jxb/erv204>
- Simkin AJ, McAusland L, Lawson T, Raines CA. Overexpression of the RieskeFeS protein increases electron transport rates and biomass yield. *Plant Physiol*. 2017b;175(1):134–145. <https://doi.org/10.1104/pp.17.00622>
- Slama V, Cupellini L, Mascoli V, Liguori N, Croce R, Mennucci B. Origin of low-lying red states in the Lhca4 light-harvesting complex of photosystem I. *J Phys Chem Lett*. 2023;14(37):8345–8352. <https://doi.org/10.1021/acs.jpcclett.3c02091>
- Slattery RA, Ort DR. Perspectives on improving light distribution and light use efficiency in crop canopies. *Plant Physiol*. 2021;185(1):34–48. <https://doi.org/10.1093/plphys/kiaa006>
- Slattery RA, VanLoocke A, Bernacchi CJ, Zhu XG, Ort DR. Photosynthesis, light use efficiency, and yield of reduced-chlorophyll soybean mutants in field conditions. *Front Plant Sci*. 2017;8:549. <https://doi.org/10.3389/fpls.2017.00549>
- Slattery RA, Walker BJ, Weber APM, Ort DR. The impacts of fluctuating light on crop performance. *Plant Physiol*. 2018;176(2):990–1003. <https://doi.org/10.1104/pp.17.01234>
- Smith EN, van Aalst M, Tosens T, Niinemets U, Stich B, Morosinotto T, Alboresi A, Erb TJ, Gomez-Coronado PA, Tolleter D, et al. Improving photosynthetic efficiency toward food security: strategies, advances, and perspectives. *Mol Plant*. 2023;16(10):1547–1563. <https://doi.org/10.1016/j.molp.2023.08.017>
- Song Q, Wang Y, Qu M, Ort DR, Zhu XG. The impact of modifying photosystem antenna size on canopy photosynthetic efficiency—development of a new canopy photosynthesis model scaling from metabolism to canopy level processes. *Plant Cell Environ*. 2017;40(12):2946–2957. <https://doi.org/10.1111/pce.13041>
- Song Q-F, Xiao H, Xiao X, Zhu XG. A new canopy photosynthesis and transpiration measurement system (CAPTS) for canopy gas exchange research. *Agr For Meteorol*. 2016;217:101–107. <https://doi.org/10.1016/j.agrformet.2015.11.020>
- Song Q-F, Zhang G, Zhu X-G. Optimal crop canopy architecture to maximise canopy photosynthetic CO<sub>2</sub> uptake under elevated CO<sub>2</sub> – a theoretical study using a mechanistic model of canopy photosynthesis. *Funct Plant Biol*. 2013;40(2):108–124. <https://doi.org/10.1071/FP12056>
- South PF, Cavanagh AP, Liu HW, Ort DR. Synthetic glycolate metabolism pathways stimulate crop growth and productivity in the field. *Science*. 2019;363(6422):eaat9077. <https://doi.org/10.1126/science.aat9077>
- Sparrow-Muñoz I, Chen TC, Burgess SJ. Recent developments in the engineering of Rubisco activase for enhanced crop yield. *Biochem Soc Trans*. 2023;51(2):627–637. <https://doi.org/10.1042/BST20221281>
- Srinivasan V, Kumar P, Long SP. Decreasing, not increasing, leaf area will raise crop yields under global atmospheric change. *Glob Chang Biol*. 2017;23(4):1626–1635. <https://doi.org/10.1111/gcb.13526>
- Stitt M, Schulze D. Does Rubisco control the rate of photosynthesis and plant-growth - an exercise in molecular ecophysiology. *Plant Cell Environ*. 1994;17(5):465–487. <https://doi.org/10.1111/j.1365-3040.1994.tb00144.x>
- Suzuki Y, Wada S, Kondo E, Yamori W, Makino A. Effects of co-overproduction of sedoheptulose-1,7-bisphosphatase and Rubisco on photosynthesis in rice. *Soil Sci Plant Nutr*. 2019;65(1):36–40. <https://doi.org/10.1080/00380768.2018.1530053>
- Szurman-Zubrzycka ME, Zbieszczak J, Marzec M, Jelonek J, Chmielewska B, Kurowska MM, Szarejko I. HorTILLUS—a rich and renewable source of induced mutations for forward/reverse genetics and pre-breeding programs in barley (*Hordeum vulgare* L.). *Front Plant Sci*. 2018;9:1–16. <https://doi.org/10.3389/fpls.2018.00216>
- Tamoi M, Nagaoko M, Myagawa Y, Shigeoki S. Contribution of fructose-1,6-bisphosphatase and sedoheptulose-1,7-bisphosphatase to the photosynthetic rate and carbon flow in the Calvin cycle in transgenic plants. *Plant Cell Physiol*. 2006;47(3):380–390. <https://doi.org/10.1093/pccp/pcj004>
- Tanaka Y, Sugano SS, Shimada T, Hara-Nishimura I. Enhancement of leaf photosynthetic capacity through increased stomatal density in Arabidopsis. *New Phytol*. 2013;198(3):757–764. <https://doi.org/10.1111/nph.12186>
- Tardy F, Créach A, Havaux M. Photosynthetic pigment concentration, organization and interconversions in a pale green Syrian landrace of barley (*Hordeum Vulgare* L., Tadmor) adapted to harsh climatic conditions. *Plant Cell Environ*. 1998;21(5):479–489. <https://doi.org/10.1046/j.1365-3040.1998.00293.x>
- Taylor G, Garassino F, Aarts MGM, Harbinson J. Improving C<sub>3</sub> photosynthesis by exploiting natural genetic variation: *Hirschfeldia incana* as a model species. *Food Energy Secur*. 2022;12(1):e420. <https://doi.org/10.1002/fes3.420>
- Tcherkez GG, Farquhar GD, Andrews TJ. Despite slow catalysis and confused substrate specificity, all ribulose bisphosphate carboxylases may be nearly perfectly optimized. *Proc Natl Acad Sci U S A*. 2006;103(19):7246–7251. <https://doi.org/10.1073/pnas.0600605103>
- Theeuwens TPJM, Logie LL, Harbinson J, Aarts MGM. Genetics as a key to improving crop photosynthesis. *J Exp Bot*. 2022;73(10):3122–3137. <https://doi.org/10.1093/jxb/erac076>
- Tholen D, Ethier G, Genty B, Pepin ZX-G. Variable mesophyll conductance revisited: theoretical background and experimental implications. *Plant Cell Environ*. 2012;35(12):2087–2103. <https://doi.org/10.1111/j.1365-3040.2012.02538.x>
- Tholen D, Zhu X-G. The mechanistic basis of internal conductance: a theoretical analysis of mesophyll cell photosynthesis and CO<sub>2</sub> diffusion. *Plant Physiol*. 2011;156(1):90–105. <https://doi.org/10.1104/pp.111.172346>
- Tikhonov AN. The cytochrome b6f complex at the crossroad of photosynthetic electron transport pathways. *Plant Physiol Biochem*. 2014;81:163–183. <https://doi.org/10.1016/j.plaphy.2013.12.011>
- Tros M, Bersanini L, Shen G, Ho MY, van Stokkum IHM, Bryant DA, Croce R. Harvesting far-red light: functional integration of chlorophyll f into Photosystem I complexes of *Synechococcus* sp. PCC 7002. *Biochim Biophys Acta*. 2020;1861(8):148206. <https://doi.org/10.1016/j.bbabi.2020.148206>
- Tros M, Mascoli V, Shen GZ, Ho MY, Bersanini L, Gisriel CJ, Bryant DA, Croce R. Breaking the red limit: efficient trapping of long-wavelength excitations in chlorophyll-f-containing photosystem I. *Chem*. 2021;7(1):155–173. <https://doi.org/10.1016/j.chempr.2020.10.024>

- Trudeau DL, Edlich-Muth C, Zarzycki J, Scheffen M, Goldsmith M, Khersonsky O, Avizemer Z, Fleishman SJ, Cotton CAR, Erb TJ, et al. Design and in vitro realization of carbon-conserving photorespiration. *Proc Natl Acad Sci U S A*. 2018;115(49):E11455–E11464. <https://doi.org/10.1073/pnas.1812605115>
- Uematsu K, Suzuki N, Iwamae T, Inui M, Yukawa H. Increased fructose 1,6-bisphosphate aldolase in plastids enhances growth and photosynthesis of tobacco plants. *J Exp Bot*. 2012;63(8):3001–3009. <https://doi.org/10.1093/jxb/ers004>
- van Bel AJE, Gamalei YV. Ecophysiology of phloem loading in source leaves. *Plant Cell Environ*. 1992;15(3):265–270. <https://doi.org/10.1111/j.1365-3040.1992.tb00973.x>
- van Bezouw R, Keurentjes JJB, Harbinson J, Aarts MGM. Converging phenomics and genomics to study natural variation in plant photosynthetic efficiency. *Plant J*. 2019;97(1):112–133. <https://doi.org/10.1111/tpj.14190>
- Viola S, Roseby W, Santabarbara S, Nurnberg D, Assuncao R, Dau H, Selles J, Boussac A, Fantuzzi A, Rutherford AW. Impact of energy limitations on function and resilience in long-wavelength Photosystem II. *Elife*. 2022;11:e79890. <https://doi.org/10.7554/eLife.79890>
- Von Caemmerer S. *Biochemical models of leaf photosynthesis*. Collingwood: CSIRO Publishing; 2000
- Von Caemmerer S, Furbank RT. Modelling C4 photosynthesis. In: **Sage RF, Monson RK**, editors. *The biology of C4 photosynthesis*. London: Academic Press; 1999. p. 173–211.
- von Caemmerer S, Furbank RT. Strategies for improving C4 photosynthesis. *Curr Opin Plant Biol*. 2016;31:125–134. <https://doi.org/10.1016/j.pbi.2016.04.003>
- Walker BJ, Drewry DT, Slaterry RA, VanLoocke A, Cho YB, Ort DR. Chlorophyll can be reduced in crop canopies with little penalty to photosynthesis. *Plant Physiol*. 2018;176(2):1215–1232. <https://doi.org/10.1104/pp.17.01401>
- Walker BJ, Vanlooche A, Bernacchi CJ, Ort DR. The costs of photorespiration to food production now and in the future. *Annu Rev Plant Biol*. 2016;67(1):107–129. <https://doi.org/10.1146/annurev-arplant-043015-111709>
- Wall S, Violet-Chabrand S, Davey P, van Rie J, Galle A, Cockram J, Lawson T. Stomata on the abaxial and adaxial leaf surface contribute differently to leaf gas exchange and photosynthesis in wheat. *New Phytol*. 2022;235(5):1743–1756. <https://doi.org/10.1111/nph.18257>
- Wang Y, Bräutigam A, Weber APM, Zhu X-G. Three distinct biochemical subtypes of C4 photosynthesis? A modelling analysis. *J Exp Bot*. 2014a;65(13):3567–3578. <https://doi.org/10.1093/jxb/eru058>
- Wang Y, Burgess SJ, de Becker EM, Long SP. Photosynthesis in the fleeting shadows: an overlooked opportunity for increasing crop productivity? *Plant J*. 2020;101(4):874–884. <https://doi.org/10.1111/tpj.14663>
- Wang P, Khoshravesh R, Karki S, Tapia R, Balahadia CP, Bandyopadhyay A, Quick WP, Furbank R, Sage TL, Langdale JA. Re-creation of a key step in the evolutionary switch from C3 to C4 leaf anatomy. *Curr Biol*. 2017;27(21):3278–3287.e6. <https://doi.org/10.1016/j.cub.2017.09.040>
- Wang P, Liang FC, Wittmann D, Siegel A, Shan S, Grimm B. Chloroplast SRP<sub>43</sub> acts as a chaperone for glutamyl-tRNA reductase, the rate-limiting enzyme in tetrapyrrole biosynthesis. *Proc Natl Acad Sci U S A*. 2018;115(15):E3588–E3596. <https://doi.org/10.1073/pnas.1719645115>
- Wang Y, Long SP, Zhu XG. Elements required for an efficient NADP-malic enzyme type C<sub>4</sub> photosynthesis. *Plant Physiol*. 2014b;164(4):2231–2246. <https://doi.org/10.1104/pp.113.230284>
- Wang Y, Smith JAC, Zhu XG, Long SP. Rethinking the potential productivity of crassulacean acid metabolism by integrating metabolic dynamics with shoot architecture, using the example of *Agave tequilana*. *New Phytol*. 2023;239(6):2180–2196. <https://doi.org/10.1111/nph.19128>
- Wang Z, Zhang T, Xing Y, Zeng X, Wang L, Liu Z, Shi J, Zhu X, Ma L, Li Y, et al. YGL9, encoding the putative chloroplast signal recognition particle 43 kDa protein in rice, is involved in chloroplast development. *J Integr Agric*. 2016;15(5):944–953. [https://doi.org/10.1016/S2095-3119\(15\)61310-7](https://doi.org/10.1016/S2095-3119(15)61310-7)
- Watanabe N, Nakada E. Seasonal variation of leaf colour in Syrian Barley and its association with photosynthetic electron transport rate. *Cereal Res Commun*. 1999;27(1–2):171–178. <https://doi.org/10.1007/BF03543934>
- Werk KS, Ehleringer J, Forseth IN, Cook CS. Photosynthetic characteristics of Sonoran Desert winter annuals. *Oecologia*. 1983;59(1):101–105. <https://doi.org/10.1007/BF00388081>
- Wong SC, Cowan IR, Farquhar GD. Stomatal conductance correlates with photosynthetic capacity. *Nature*. 1979;282(5737):424–426. <https://doi.org/10.1038/282424a0>
- Wu A, Brider J, Busch FA, Chen M, Chenu K, Clarke VC, Collins B, Ermakova M, Evans JR, Farquhar GD, et al. A cross-scale analysis to understand and quantify the effects of photosynthetic enhancement on crop growth and yield across environments. *Plant Cell Environ*. 2023;46(1):23–44. <https://doi.org/10.1111/pce.14453>
- Xiao Y, Sloan J, Hepworth C, Fradera-Soler M, Mathers A, Thorley R, Baillie A, Jones H, Chang T, Chen X, et al. Defining the scope for altering rice leaf anatomy to improve photosynthesis: a modelling approach. *New Phytol*. 2022;237(2):441–453. <https://doi.org/10.1111/nph.18564>
- Xiao Y, Tholen D, Zhu X-G. The influence of leaf anatomy on the internal light environment and photosynthetic electron transport rate: exploration with a new leaf ray tracing model. *J Exp Bot*. 2016;67(21):6021–6035. <https://doi.org/10.1093/jxb/erw359>
- Xiao Y, Zhu X-G. Components of mesophyll resistance and their environmental responses: a theoretical modelling analysis. *Plant Cell Environ*. 2017;40(11):2729–2742. <https://doi.org/10.1111/pce.13040>
- Xin CP, Tholen D, Devloo V, Zhu XG. The benefits of photorespiratory bypasses: how can they work? *Plant Physiol*. 2015;167(2):574–585. <https://doi.org/10.1104/pp.114.248013>
- Yamamoto HY, Nakayama TOM, Chichester CO. Studies on the light and dark interconversions of leaf xanthophylls. *Arch Biochem Biophys*. 1962;97(1):168–173. [https://doi.org/10.1016/0003-9861\(62\)90060-7](https://doi.org/10.1016/0003-9861(62)90060-7)
- Yamori W, Hikosaka K, Way DA. Temperature response of photosynthesis in C3, C4, and CAM plants: temperature acclimation and temperature adaptation. *Photosynth Res*. 2014;119(1–2):101–117. <https://doi.org/10.1007/s11120-013-9874-6>
- Yamori W, Takahashi S, Makino A, Price GD, Badger MR, Von Caemmerer S. The roles of ATP synthase and the cytochrome b6f complexes in limiting chloroplast electron transport and determining photosynthetic capacity. *Plant Physiol*. 2011;155(2):956–962. <https://doi.org/10.1104/pp.110.168435>
- Yi XP, Yao HS, Fan DY, Zhu XG, Losciale P, Zhang Y, Zhang W, Chow FWS. The energy cost of repairing photoinactivated photosystem II: an experimental determination in cotton leaf discs. *New Phytol*. 2022;235(2):446–456. <https://doi.org/10.1111/nph.18165>
- Yin X, Tang M, Xia X, Yu J. BRASSINAZOLE RESISTANT 1 mediates brassinosteroid-induced Calvin cycle to promote photosynthesis in tomato. *Front Plant Sci*. 2022;12:811948. <https://doi.org/10.3389/fpls.2021.811948>
- Yoon DK, Ishiyama K, Suganami M, Tazoe Y, Watanabe M, Imaruoka S, Ogura M, Ishida H, Suzuki Y, Obara M, et al. Transgenic rice

- overproducing Rubisco exhibits increased yields with improved nitrogen-use efficiency in an experimental paddy field. *Nat Food*. 2020;1(2):134–139. <https://doi.org/10.1038/s43016-020-0033-x>
- Zaks J, Amarnath K, Kramer DM, Niyog KK, Fleming GR. A kinetic model of rapidly reversible nonphotochemical quenching. *Proc Natl Acad Sci U S A*. 2012;109(39):15757–15762. <https://doi.org/10.1073/pnas.1211017109>
- Zelitch I, Schultes NP, Peterson RB, Brown P, Brutnell TP. High glycolate oxidase activity is required for survival of maize in normal air. *Plant Physiol*. 2009;149(1):195–204. <https://doi.org/10.1104/pp.108.128439>
- Zhang GG, Sakai H, Tokida T, Usui Y, Zhu C, Nakamura H, Yoshimoto M, Fukuoka M, Kobayashi K, Hasegawa T. The effects of free-air CO<sub>2</sub> enrichment (FACE) on carbon and nitrogen accumulation in grains of rice (*Oryza sativa* L.). *J Exp Bot*. 2013;64(11):3179–3188. <https://doi.org/10.1093/jxb/ert154>
- Zhang DY, Sun GJ, Jiang XH. Donald's ideotype and growth redundancy: a game theoretical analysis. *Field Crops Res*. 1999;61(2):179–187. [https://doi.org/10.1016/S0378-4290\(98\)00156-7](https://doi.org/10.1016/S0378-4290(98)00156-7)
- Zhou Z, Struik PC, Gu J, van der Putten PE, Wang Z, Yin X, Yang J. Enhancing leaf photosynthesis from altered chlorophyll content requires optimal partitioning of nitrogen. *Crop Environ*. 2023;2(1):24–36. <https://doi.org/10.1016/j.crope.2023.02.001>
- Zhu X-G, De Sturler E, Long SP. Optimizing the distribution of resources between enzymes of carbon metabolism can dramatically increase photosynthetic rate: a numerical simulation using an evolutionary algorithm. *Plant Physiol*. 2007;145(2):513–526. <https://doi.org/10.1104/pp.107.103713>
- Zhu X-G, Long SP, Ort DR. Improving photosynthetic efficiency for greater yield. *Ann Rev Plant Biol*. 2010;61(1):235–261. <https://doi.org/10.1146/annurev-arplant-042809-112206>
- Zhu X-G, Ort DR, Whitmarsh J, Long SP. The slow reversibility of photosystem II thermal energy dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies: a theoretical analysis. *J Exp Bot*. 2004;55(400):1167–1175. <https://doi.org/10.1093/jxb/erh141>
- Zhu X-G, Song Q-F, Ort DR. Elements of a dynamic systems model of canopy photosynthesis. *Curr Opin Plant Biol*. 2012;15(3):237–244. <https://doi.org/10.1016/j.pbi.2012.01.010>
- Zhu X-G, Wang Y, Ort DR, Long SP. e-Photosynthesis: a comprehensive dynamic mechanistic model of C3 photosynthesis: from light capture to sucrose synthesis. *Plant Cell Environ*. 2013;36(9):1711–1727. <https://doi.org/10.1111/pce.12025>